

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 5 : C07H 21/00, C12Q 1/68</p>	<p>A1</p>	<p>(11) International Publication Number: WO 91/17174 (43) International Publication Date: 14 November 1991 (14.11.91)</p>
<p>(21) International Application Number: PCT/US91/03200 (22) International Filing Date: 8 May 1991 (08.05.91) (30) Priority data: 520,716 8 May 1990 (08.05.90) US (71) Applicant (for all designated States except US): NEUROGENETIC CORPORATION [US/US]; 215 College Road, Pacamus, NJ 07652 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WEINSHANK, Richard, L. [US/US]; 302 West 87th Street, Apt. 34, New York, NY 10024 (US). BRANCHEK, Theresa [US/US]; 541 Martence Avenue, Teaneck, NJ 07766 (US). HARTIG, Paul, R. [US/US]; 179 Linden Street, Mahwah, NJ 07430 (US).</p>		<p>(74) Agent: WHITE, John, P.; Cooper & Dunham, 30 Rockefeller Plaza, New York, NY 10112 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), SU, US. Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>

(54) Title: DNA ENCODING HUMAN 5-HT_{1D} RECEPTORS AND USES THEREOF

(57) Abstract

This invention provides isolated nucleic acid molecules encoding human 5-HT_{1D} receptors, isolated proteins which are human 5-HT_{1D} receptors, vectors comprising isolated nucleic acid molecules encoding human 5-HT_{1D} receptors, mammalian cells comprising such vectors, antibodies directed to the human 5-HT_{1D} receptors, nucleic acid probes useful for detecting nucleic acid encoding human 5-HT_{1D} receptors, antisense oligonucleotides complementary to any sequences of a nucleic acid molecule which encodes a human 5-HT_{1D} receptor, pharmaceutical compounds related to human 5-HT_{1D} receptors, and nonhuman transgenic animals which express DNA a normal or a mutant human 5-HT_{1D} receptor. This invention further provides methods for determining ligand binding, detecting expression, drug screening, and treatment involving the human 5-HT_{1D} receptor.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland			SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TG	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		

DNA ENCODING HUMAN 5-HT₁₀ RECEPTORS AND USES THEREOF

This application is a continuation-in-part of U.S. Serial No. 520,716, filed May 8, 1990, the contents of which are hereby incorporated by reference into the subject application.

5

Background of the Invention

10

Throughout this application various publications are referenced by full citations within parentheses. The disclosures of these publications in their entireties are hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains.

15

20

Pharmacological studies, and more recently gene cloning, have established that multiple receptor subtypes exist for most, if not all, neurotransmitters. The existence of multiple receptor subtypes provides one mechanism by which a single neurotransmitter can elicit distinct cellular responses. The variation in cellular response can be achieved by the association of individual receptor subtypes with different G proteins and different signalling systems. Further flexibility is provided by the ability of distinct receptors for the same ligand to activate or inhibit the same second messenger system.

25

30

Individual receptor subtypes reveal characteristic differences in their abilities to bind a number of ligands, but the structural basis for the distinct ligand-binding properties is not known. Physiologists and pharmacologists have attempted to specify particular biological functions or anatomical locations for some receptor subtypes, but this

35

-2-

has met with limited success. Similarly, the biochemical mechanisms by which these receptors transduce signals across the cell surface have been difficult to ascertain without having well-defined cell populations which express exclusively one receptor subtype.

5

While all the receptors of the serotonin type recognize serotonin, several pharmacologically distinct subtypes of serotonin receptors have been identified, and given a classification name 5-HT_x, where X identifies the subtype. In many cases, these subtypes have been or will be associated with single gene products, but in some cases a single subtype may be found to contain several different receptor proteins (gene products) or two different subtypes may be later shown to arise from different properties of the same receptor protein which are exhibited when it is expressed in different tissue environments. In many cases, different serotonin receptor subtypes have been shown to couple to different second messenger pathways that are linked through guanine-nucleotide regulatory (G) proteins.

20

Radioligand filtration binding techniques have been employed for over ten years in an effort to more completely characterize receptor subtypes within the serotonin receptor family (Schmidt and Peroutka, FASEB J. 3:2242 (1989)). Using these methods, two broad classes of G protein-coupled serotonin receptors have been described, 5-HT₁, and 5-HT₂. These differ in their selectivity for drugs. 5-HT₁ receptors display high (nanomolar) affinity for serotonin and can be labeled with [³H]5-HT. 5-HT₂ receptors display low affinity for serotonin but have high (nanomolar) affinity for antagonists such as Ketanserin, Mesulergine,

30

35

-3-

Metergoline and d-LSD.

Within the 5-HT₁ receptor class, several subtypes have been distinguished on the basis of their pharmacological binding profiles, second messenger coupling and physiological roles. One such subtype, the 5-HT₁₀ receptor, was originally defined as a particular type of [³H]5-HT binding site in the bovine caudate (Heuring and Peroutka, J. Neurosci. 7:894 (1987)). This definition was not based on properties of a single purified receptor protein or single gene product, but rather was based on experimental observations in a model tissue. As discussed below, later research has shown that there may be multiple receptor proteins (known as subtypes) within this model tissue, all of which contribute to the binding profile that was used to define the 5-HT₁₀ receptor.

The 5-HT₁₀ receptor subtype has been shown to inhibit adenylate cyclase activity (Schoeffter, P. and Hoyer, D., Naunyn-Schmiedeberg's Arch. Pharmacol. 340:285 (1989)). The 5-HT₁₀ receptor subtype has also been characterized in guinea pig (Waeber, et al. Naunyn-Schmiedeberg's Arch. Pharmacol. 340:479-485 (1989)), pigeon (Waeber, 1989), pig (Waeber, et al. Naunyn-Schmiedeberg's Arch. Pharmacol. 377:595-601 (1988)), calf (Waeber, et al. (1988)) and human brain (Waeber, et al. (1988); Herrick-Davis and Titeler, J. Neurochem, 50:1624-1631 (1988)). Among the other serotonin receptor subtypes, the 5-HT_{1A}, and 5-HT_{1B} receptors inhibit adenylate cyclase, and 5-HT_{1C} and 5-HT₂ receptors activate phospholipase C pathways, stimulating breakdown of polyphosphoinositides (Schmidt and Peroutka, FASEB J. 3:2242 (1989)).

The pharmacological actions of sumatriptan (GR43175), a new

anti-migraine medication under development by Glaxo Pharmaceutical Corp., have been linked to the 5-HT_{1D} receptor site (Peroutka and McCarthy, Eur. J. Pharmacology 163:133 (1989)); Schoeffter and Hoyer, Naunyn-Schmiedeberg Arch. Pharmacology 340:135 (1989)). Recently, one report has shown that the 5-HT_{1D} binding site of piglet caudate could be subdivided into two sites, based on the binding affinities of sumatriptan and 5-carboxamidotryptamine (5-CT) (Sumner and Humphrey, Br. J. Pharmacol. 98:29 (1989)). One of these binding sites, with low affinity for sumatriptan and 5-CT, resembles the 5-HT_{1E} site of human cortex (Leonhardt, Herrick-Davis and Titeler, J. Neurochem. 53:465 (1989)) while the binding site with high affinity for these compounds resembles the classic 5-HT_{1D} receptor, and the site of action of sumatriptan.

Another study, by Xiong and Nelson (Life Sci. 45:1433-1442 (1989)) indicated that a high affinity [³H]5-HT binding site in the rabbit caudate, termed the 5-HT_{1R} binding site, is similar to, but pharmacologically distinct from, the 5-HT_{1D} binding site described in the bovine caudate. These authors presented data indicating that two drugs, spiperone and spirilene, exhibited significantly lower affinity for the 5-HT_{1R} binding site than for the 5-HT_{1D} receptor, and noted several other differences in binding properties between these sites. Investigation of the bovine caudate in light of these findings led to the conclusion that there may be a component of the 5-HT_{1D} receptor in bovine caudate that represents a 5-HT_{1R} binding site. Alternatively, the authors speculated that the 5-HT_{1D} binding site in the bovine caudate may be a heterogenous group of sites with similar properties. As noted by the authors, it is clear that additional work will be need to clarify these issues.

-5-

A gene for a G protein-coupled receptor was recently isolated by Libert, et al. from a dog cDNA library (Science 244:569-572, 1989). This gene, termed the RDC4 gene, was not expressed by these authors, and therefore no characterization of the properties of the protein encoded by this gene was made. The dog RDC4 gene was isolated and expressed by the applicants, and was determined by the applicants to encode a 5-HT_{1D} receptor (not published).

The serotonin 5-HT_{1D} receptors belong to a family of receptors which are distinguished by their seven-transmembrane configuration and their functional linkage to G-proteins. This family includes rhodopsin and related opsins (Nathans, J. and Hogness, D.S., Cell 34:807 (1983)), the α and β adrenergic receptors (Dohlman, H.G., et al., Biochemistry 26:2657 (1987)), the muscarinic cholinergic receptors (Bonner, T.I., et al., Science 237:527 (1987)), the substance K neuropeptide receptor, (Masu, Y., et al., Nature 329:836 (1987)), the yeast mating factor receptors, (Burkholder, A.C. and Hartwell, L.H., Nucl. Acids Res. 13:8463(1985); Hagan, D.C., et al., Proc. Natl. Acad. Sci. USA 83:1418 (1986)); Nakayama, N. et al., EMBO J. 4:2643 (1985)), and the oncogene c-mas, (Young, et al., Cell 45:711 (1986)). Each of these receptors is thought to transduce extracellular signals by interaction with guanine nucleotide-binding (G) proteins (Dohlman, H.G., et al., Biochemistry 26:2657 (1987); Dohlman, H.G., et al., Biochemistry 27:1813 (1988); O'Dowd, B.F., et al., Ann.Rev. Neurosci., in press).

Summary of the Invention

This invention provides an isolated nucleic acid molecule encoding a human 5-HT_{1D} receptor.

5

This invention also provides an isolated protein which is a human 5-HT_{1D} receptor.

10

This invention provides a vector comprising an isolated nucleic acid molecule encoding a human 5-HT_{1D} receptor.

15

This invention also provides vectors such as plasmids comprising a DNA molecule encoding a human 5-HT_{1D} receptor, adapted for expression in a bacterial cell, a yeast cell, or a mammalian cell which additionally comprise the regulatory elements necessary for expression of the DNA in the bacterial, yeast, or mammalian cells so located relative to the DNA encoding a human 5-HT_{1D} receptor as to permit expression thereof.

20

This invention provides a mammalian cell comprising a DNA molecule encoding a human 5-HT_{1D} receptor.

25

This invention provides a method for determining whether a ligand not known to be capable of binding to a human 5-HT_{1D} receptor can bind to a human 5-HT_{1D} receptor which comprises contacting a mammalian cell comprising a DNA molecule encoding a human 5-HT_{1D} receptor with the ligand under conditions permitting binding of ligands known to bind to the 5-HT_{1D} receptor, detecting the presence of any of the ligand bound to the 5-HT_{1D} receptor, and thereby determining whether the ligand binds to the 5-HT_{1D} receptor.

30

35

-7-

5 This invention also provides a method of screening drugs to identify drugs which specifically interact with, and bind to, the human 5-HT_{1D} receptor on the surface of a cell which comprises contacting a mammalian cell comprising a DNA molecule encoding a human 5-HT_{1D} receptor on the surface of a cell with a plurality of drugs, determining those drugs which bind to the mammalian cell, and thereby identifying drugs which specifically interact with, and bind to, the human 5-HT_{1D} receptor.

10 This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding a human 5-HT_{1D} receptor.

15 This invention also provides a method of detecting expression of a 5-HT_{1D} receptor on the surface of a cell by detecting the presence of mRNA coding for a 5-HT_{1D} receptor which comprises obtaining total mRNA from the cell and
20 contacting the mRNA so obtained with a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding a human 5-HT_{1D} receptor under hybridizing
25 conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the 5-HT_{1D} receptor by the cell.

30 This invention provides an antisense oligonucleotide having a sequence capable of binding specifically with any sequences of an mRNA molecule which encodes a human 5-HT_{1D} receptor so as to prevent translation of the mRNA molecule.

This invention provides an antibody directed to the human 5-HT_{1D} receptor.

5 This invention provides a transgenic nonhuman mammal expressing DNA encoding a human 5-HT_{1D} receptor. This invention also provides a transgenic nonhuman mammal expressing DNA encoding a human 5-HT_{1D} receptor so mutated as to be incapable of normal receptor activity, and not expressing native 5-HT_{1D} receptor. This invention further provides a transgenic nonhuman mammal whose genome comprises
10 antisense DNA complementary to DNA encoding a human 5-HT_{1D} receptor so placed as to be transcribed into antisense mRNA which is complementary to mRNA encoding a 5-HT_{1D} receptor and which hybridizes to mRNA encoding a 5-HT_{1D} receptor thereby reducing its translation.

15 This invention provides a method of determining the physiological effects of expressing varying levels of human 5-HT_{1D} receptors which comprises producing a transgenic nonhuman animal whose levels of human 5-HT_{1D} receptor expression are varied by use of an inducible promoter which
20 regulates human 5-HT_{1D} receptor expression.

This invention also provides a method of determining the physiological effects of expressing varying levels of human 5-HT_{1D} receptors which comprises producing a panel of
25 transgenic nonhuman animals each expressing a different amount of human 5-HT_{1D} receptor.

30 This invention provides a method for diagnosing a predisposition to a disorder associated with the expression of a specific human 5-HT_{1D} receptor allele which comprises:
a. obtaining DNA of subjects suffering from the disorder;
b. performing a restriction digest of the DNA with a panel of

restriction enzymes; c.electrophoretically separating the resulting DNA fragments on a sizing gel; d.contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing to DNA encoding a human 5-HT₁₀ receptor and labelled with a detectable marker; e.detecting labelled bands which have hybridized to the the DNA encoding a human 5-HT₁₀ receptor labelled with a detectable marker to create a unique band pattern specific to the DNA of subjects suffering from the disorder; f.preparing DNA obtained for diagnosis by steps a-e; and g.comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step e and the DNA obtained for diagnosis from step f to determine whether the patterns are the same or different and thereby to diagnose predisposition to the disorder if the patterns are the same. This method may also be used to diagnose a disorder associated with the expression of a specific human 5-HT₁₀ receptor allele is diagnosed.

This invention provides a method of preparing the isolated 5-HT₁₀ receptor which comprises inducing cells to express 5-HT₁₀ receptor, recovering the receptor from the resulting cells, and purifying the receptor so recovered.

This invention provides a method of preparing the isolated 5-HT₁₀ receptor which comprises inserting nucleic acid encoding 5-HT₁₀ receptor in a suitable vector, inserting the resulting vector in a suitable host cell, recovering the receptor produced by the resulting cell, and purifying the receptor so recovered.

This invention provides an antisense oligonucleotide having a sequence capable of binding specifically with any sequences of an mRNA molecule which encodes a receptor so as

-10-

to prevent translation of the mRNA molecule.

This invention also provides a transgenic nonhuman mammal expressing DNA encoding a receptor.

5 This invention further provides a transgenic nonhuman mammal expressing DNA encoding a receptor so mutated as to be incapable of normal receptor activity, and not expressing native receptor.

10 This invention provides a method of determining the physiological effects of expressing varying levels of a receptor which comprises producing a transgenic nonhuman animal whose levels of the receptor expression are varied by use of an inducible promoter which regulates receptor
15 expression.

This invention also provides a method of determining the physiological effects of expressing varying levels of a receptor which comprises producing a panel of transgenic
20 nonhuman animals each expressing a different amount of the receptor.

This invention further provides transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA
25 encoding a receptor so placed as to be transcribed into antisense mRNA which is complementary to mRNA encoding the receptor and which hybridizes to mRNA encoding the receptor thereby preventing its translation.

30 This invention provides a method for determining whether a ligand not known to be capable of binding to a receptor can bind to a receptor which comprises contacting a mammalian

35

-11-

cell comprising an isolated DNA molecule encoding a receptor with the ligand under conditions permitting binding of ligands known to bind to a receptor, detecting the presence of any of the ligand bound to the receptor, and thereby determining whether the ligand binds to a receptor.

5

10

15

20

25

30

35

-12-

Brief Description of the Figures

Figure 1. cDNA clones representing gene 5-HT₁₀₋₁ (gene 8-30-84).

5 Two cDNA clones were isolated from a human hippocampus (hH) library by screening approximately 10⁶ recombinants in a Lambda Zap II vector. Clones hH-13 and hH-46 spanned the entire coding region of gene 5-HT₁₀₋₁. The seven presumed α -helical membrane-spanning domains (TM-1 to TM-7) are shown and are separated by extracellular (o1-o4) and intracellular (i1-i4) loops.

Figure 2. Restriction map of gene 5-HT₁₀₋₂ (gene 11).

15 Clone hP-11, containing approximately 15 kilobase (kb) of human genomic DNA was obtained by screening approximately 2 x 10⁶ recombinants from a Lambda Fix II human placental (hP) genomic library (Stratagene) with the ~1.3kb Hind III-Sph-I dog RDC4 probe. The seven presumed α -helical membrane-spanning domains (TM-1 to TM-7) of the deduced amino acid sequence are shown below and are separated by extracellular (o1-o4) and intracellular (i1-i4) loops. Restrictions sites are indicated.

25 Figure 3. Nucleotide and deduced amino acid sequence of gene 5-HT₁₀₋₁ (gene 8-30-84).

30 Numbers above the nucleotide sequence indicate nucleotide position. DNA sequence was determined by the chain termination method of Sanger, et al., on denatured doubled-stranded plasmid templates using the enzyme Sequenase. Deduced amino acid sequence (single letter code) of a long

35

-13-

open reading frame is shown. A plasmid comprising the cDNA sequence shown in Figure 3 has been constructed and is designated pcEXV-8-30-84. The plasmid pcEXV-8-30-84 was deposited on April 17, 1990 pursuant to the provisions of the Budapest Treaty On The International Recognition Of The Deposit Of Microorganisms For The Purposes Of Patent Procedure with the Patent Culture Depository of the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland, 20852, U.S.A. under ATCC Accession No. 40790.

Figure 4. Nucleotide and deduced amino acid sequence of gene 5-HT₁₀₋₂ (gene 11).

Numbers above the nucleotide sequence indicate nucleotide position. DNA sequence was determined by the chain termination method of Sanger, et al., on denatured double-stranded plasmid templates using the enzyme Sequenase. Deduced amino acid sequence (single letter code) of a long open reading frame is shown. A plasmid comprising the cDNA sequence shown in Figure 4 has been constructed and is designated pSVL-11. The plasmid pSVL-11 was deposited on April 17, 1990 pursuant to the provisions of the Budapest Treaty On The International Recognition Of The Deposit Of Microorganisms For The Purposes Of Patent Procedure with the Patent Culture Depository of the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland, 20852, U.S.A. under ATCC Accession No. 40791.

Figure 5. Seven transmembrane-spanning model of the deduced amino acid sequence of gene 5-HT₁₀₋₁ (Peptide 8-30-84).

Amino acids in shaded circles are amino acids which differ

-14-

from the amino acids at the corresponding position in the dog RDC4 receptor. The arrows indicate potential sites of N-linked glycosylation. Amino acids in circles that straddle the lines which represent the membrane are considered outside the transmembrane region for purposes of this application.

Figure 6. Comparison of the human 5-HT₁₀ receptor primary structures with other serotonin receptors.

Amino acid sequences (single letter code) are aligned to optimize homology. The putative transmembrane spanning domains are indicated by stars and identified by Roman numerals (TM I-VII).

-15-

Detailed Description of the Invention

As used herein, the 5-HT receptor family is defined as the group of mammalian proteins that function as receptors for serotonin. A 5-HT receptor subfamily is defined as a subset of proteins belonging to the 5-HT receptor family which are encoded by genes which exhibit homology of 65% or higher with each other in their deduced amino acid sequences within presumed transmembrane regions (linearly contiguous stretches of hydrophobic amino acids, bordered by charged or polar amino acids, that are long enough to form secondary protein structures that span a lipid bilayer). Three human 5-HT receptor subfamilies can be distinguished based on the information presently available. The 5-HT₂ receptor subfamily contains the human 5-HT₂ receptor. Although no other human members of this family have been described, the rat 5-HT₂ receptor (Pritchett, et al. 1988; Julius, et al. Proc. Natl. Acad. Sci. USA 87:928-932, 1990) and the rat 5HT_{1c} receptor (Julius, et al. 1988) constitute a rat 5-HT receptor subfamily. The 5-HT_{1A} subfamily contains the human 5-HT_{1A} receptor, also known as G-21 (Fargin, et al. 1988). The 5-HT₁₀ receptor subfamily contains two members, the 5-HT₁₀₋₁ receptor (also termed 5-HT_{10a} and/or peptide 8-30-84) and the 5-HT₁₀₋₂ receptor (also termed 5-HT_{10b} and/or peptide 11) which are described herein. Therefore, the term "human 5-HT₁₀ receptor" as used herein is defined as meaning a member of the 5-HT₁₀ receptor subfamily described above. Although this definition differs from the pharmacological definition used earlier, there is significant overlap between the present definition and the pharmacological definition. Members of the 5-HT₁₀ receptor subfamily so described include the 5-HT₁₀₋₁ receptor, the 5-HT₁₀₋₂ receptor, and any other receptors which have a 65% homology to the DNA

-16-

and amino acid sequence shown in Figure 3 or to the DNA and amino acid sequence shown in Figure 4, according to the definition of "subfamily". This invention relates to the discovery of the first members of the human 5-HT₁₀ receptor subfamily.

5 This invention provides an isolated nucleic acid molecule encoding a human 5-HT₁₀ receptor. Such a receptor is by definition a member of the 5-HT₁₀ receptor subfamily. Therefore, any receptor which meets the defining criteria
10 given above is a human 5-HT₁₀ receptor. One means of isolating a human 5-HT₁₀ receptor is to probe a human genomic library with a natural or artificially designed DNA probe, using methods well known in the art. DNA probes derived from the human receptor genes 5-HT₁₀₋₁ and 5-HT₁₀₋₂ are
15 particularly useful probes for this purpose. DNA and cDNA molecules which encode human 5-HT₁₀ receptors may be used to obtain complementary genomic DNA, cDNA or RNA from human, mammalian or other animal sources, or to isolate related cDNA or genomic clones by the screening of cDNA or genomic
20 libraries, by methods described in more detail below. Transcriptional regulatory elements from the 5' untranslated region of the isolated clones, and other stability, processing, transcription, translation, and tissue specificity-determining regions from the 3' and 5'
25 untranslated regions of the isolated genes are thereby obtained. Examples of a nucleic acid molecule are an RNA, cDNA, or isolated genomic DNA molecule encoding a human 5-HT₁₀ receptor. Such molecules may have coding sequences substantially the same as the coding sequence shown in
30 Figure 3 or the coding sequence shown in Figure 4, or may have coding sequences that are 65% or more homologous to the coding sequence shown in Figure 3 or the coding sequence

-17-

shown in Figure 4. The DNA molecules of Figure 3 and Figure 4 encode the sequences of the human 5-HT₁₀ receptor genes 5-HT₁₀₋₁ and 5-HT₁₀₋₂.

5 This invention further provides a cDNA molecule of encoding a human 5-HT₁₀₋₁ receptor having a coding sequence substantially the same as the coding sequence shown in Figure 3, and a cDNA molecule encoding a human 5-HT₁₀₋₂ receptor having a coding sequence substantially the same as the coding sequence shown in Figure 4. These molecules are
10 obtained by the means described above.

This invention also provides an isolated protein which is a human 5-HT₁₀ receptor. Examples of such proteins are an isolated protein having substantially the same amino acid
15 sequence as the amino acid sequence shown in Figure 3 and an isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 4, which are human 5-HT₁₀ receptors 5-HT₁₀₋₁ and 5-HT₁₀₋₂. One means for obtaining isolated 5-HT₁₀ receptor is to express DNA encoding the receptor in a suitable host, such as a bacterial, yeast,
20 or mammalian cell, using methods well known in the art, and recovering the receptor protein after it has been expressed in such a host, again using methods well known in the art. The receptor may also be isolated from cells which express it, in particular from cells which have been transfected
25 with the expression vectors described below in more detail.

This invention provides a vector comprising an isolated nucleic acid molecule such as DNA, RNA, or cDNA encoding a
30 human 5-HT₁₀ receptor. Examples of vectors are viruses such as bacteriophages (such as phage lambda), cosmids, plasmids (such as pUC18, available from Pharmacia, Piscataway, NJ),

-18-

and other recombination vectors. Nucleic acid molecules are inserted into vector genomes by methods well known in the art. For example, insert and vector DNA can both be exposed to a restriction enzyme to create complementary ends on both molecules which base pair with each other and are then ligated together with a ligase. Alternatively, linkers can be ligated to the insert DNA which correspond to a restriction site in the vector DNA, which is then digested with the restriction enzyme which cuts at that site. Other means are also available. Specific examples of plasmids are a plasmid comprising cDNA having a coding sequence substantially the same as the coding sequence shown in Figure 3 and designated clone p5HT-8-30-84, and a plasmid comprising cDNA having a coding sequence substantially the same as the coding sequence shown in Figure 4 and designated clone p5HT-11.

This invention also provides vectors comprising a DNA molecule encoding a human 5-HT₁₀ receptor, adapted for expression in a bacterial cell, a yeast cell, or a mammalian cell which additionally comprise the regulatory elements necessary for expression of the DNA in the bacterial, yeast, or mammalian cells so located relative to the DNA encoding a human 5-HT₁₀ receptor as to permit expression thereof. DNA having coding sequences substantially the same as the coding sequence shown in Figure 3 or the coding sequence shown in Figure 4 may usefully be inserted into the vectors to express human 5-HT₁₀ receptors. Regulatory elements required for expression include promoter sequences to bind RNA polymerase and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector includes a promoter such as the lac promoter and for transcription initiation the Shine-Dalgarno sequence and the

-19-

start codon AUG (Maniatis, et al., Molecular Cloning, Cold Spring Harbor Laboratory, 1982). Similarly, a eukaryotic expression vector includes a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome. Such vectors may be obtained commercially or assembled from the sequences described by methods well known in the art, for example the methods described above for constructing vectors in general. Expression vectors are useful to produce cells that express the receptor. Certain uses for such cells are described in more detail below.

This invention further provides a plasmid adapted for expression in a bacterial, yeast, or, in particular, a mammalian cell which comprises a DNA molecule encoding a human 5-HT_{1D} receptor and the regulatory elements necessary for expression of the DNA in the bacterial, yeast, or mammalian cell so located relative to the DNA encoding a human 5-HT_{1D} receptor as to permit expression thereof. Some plasmids adapted for expression in a mammalian cell are pSVL (available from Pharmacia, Piscataway, NJ) and pcEXV-3 (Miller J. and Germain R.N., J. Exp. Med. 164:1478 (1986)). Specific examples of such plasmids are a plasmid adapted for expression in a mammalian cell comprising cDNA having coding sequences substantially the same as the coding sequence shown in Figure 3 and the regulatory elements necessary for expression of the DNA in the mammalian cell which is designated pcEXV-8-30-84 and deposited under ATCC Accession No. 40790, and a plasmid adapted for expression in a mammalian cell comprising cDNA having coding sequences substantially the same as the coding sequence shown in Figure 4 and the regulatory elements necessary for

-20-

expression of the DNA in the mammalian cell which is designated pSVL-11 and deposited under ATCC Accession No. 40791. Those skilled in the art will readily appreciate that numerous plasmids adapted for expression in a mammalian cell which comprise DNA of encoding human 5-HT₁₀ receptors and the regulatory elements necessary to express such DNA in the mammalian cell may be constructed utilizing existing plasmids and adapted as appropriate to contain the regulatory elements necessary to express the DNA in the mammalian cell. The plasmids may be constructed by the methods described above for expression vectors and vectors in general, and by other methods well known in the art.

These deposits discussed supra, and the other deposits discussed herein, were made pursuant to, and in satisfaction of, the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure with the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland 20852.

This invention provides a mammalian cell comprising a DNA molecule encoding a human 5-HT₁₀ receptor, such as a mammalian cell comprising a plasmid adapted for expression in a mammalian cell, which comprises a DNA molecule encoding a human 5-HT₁₀ receptor and the regulatory elements necessary for expression of the DNA in the mammalian cell so located relative to the DNA encoding a human 5-HT₁₀ receptor as to permit expression thereof. Numerous mammalian cells may be used as hosts, including, for example, the mouse fibroblast cell NIH3T3, CHO cells, HeLa cells, Ltk- cells, etc. A particular example of an Ltk- cell is a cell designated Ltk-8-30-84 and deposited under ATCC Accession No. CRL 10421, and comprises the plasmid designated pcEXV-8-30-84. Another

-21-

example is a cell designated Ltk-11 and deposited under ATCC Accession No. CRL 10422, and which comprises the plasmid designated pSVL-11. Expression plasmids such as those described supra may be used to transfect mammalian cells by methods well known in the art such as calcium phosphate precipitation, or DNA encoding these 5-HT₁₀ receptors may be otherwise introduced into mammalian cells, e.g., by microinjection, to obtain mammalian cells which comprise DNA, e.g., cDNA or a plasmid, encoding either human 5-HT₁₀ receptor.

This invention provides a method for determining whether a ligand not known to be capable of binding to a human 5-HT₁₀ receptor can bind to a human 5-HT₁₀ receptor which comprises contacting a mammalian cell comprising a DNA molecule encoding a human 5-HT₁₀ receptor with the ligand under conditions permitting binding of ligands known to bind to the 5-HT₁₀ receptor, detecting the presence of any of the ligand bound to the 5-HT₁₀ receptor, and thereby determining whether the ligand binds to the 5-HT₁₀ receptor. The DNA in the cell may have a coding sequence substantially the same as the coding sequence shown in Figure 3, or the coding sequence shown in Figure 4. Preferably, the mammalian cell is nonneuronal in origin. An example of a nonneuronal mammalian cell is an Ltk- cell, in particular the Ltk- cell designated L-5HT-8-30-84, or the Ltk- cell designated L-5HT-11. The preferred method for determining whether a ligand is capable of binding to the human 5-HT₁₀ receptors comprises contacting a transfected nonneuronal mammalian cell (i.e. a cell that does not naturally express any type of 5-HT or G-protein coupled receptor, thus will only express such a receptor if it is transfected into the cell) expressing a 5-HT₁₀ receptor on its surface, or contacting a membrane

-22-

preparation derived from such a transfected cell, with the ligand under conditions which are known to prevail, and thus to be associated with, in vivo binding of the ligands to a 5-HT₁₀ receptor, detecting the presence of any of the ligand being tested bound to the 5-HT₁₀ receptor on the surface of the cell, and thereby determining whether the ligand binds to the 5-HT₁₀ receptor. This response system is obtained by transfection of isolated DNA into a suitable host cell containing the desired second messenger system such as phosphoinositide hydrolysis, adenylate cyclase, guanylate cyclase or ion channels. Such a host system is isolated from pre-existing cell lines, or can be generated by inserting appropriate components of second messenger systems into existing cell lines. Such a transfection system provides a complete response system for investigation or assay of the activity of human 5-HT₁₀ receptors with ligands as described above. Transfection systems are useful as living cell cultures for competitive binding assays between known or candidate drugs and ligands which bind to the receptor and which are labeled by radioactive, spectroscopic or other reagents. Membrane preparations containing the receptor isolated from transfected cells are also useful for these competitive binding assays. Functional assays of second messenger systems or their sequelae in transfection systems act as assays for binding affinity and efficacy in the activation of receptor function. A transfection system constitutes a "drug discovery system" useful for the identification of natural or synthetic compounds with potential for drug development that can be further modified or used directly as therapeutic compounds to activate or inhibit the natural functions of the human 5-HT₁₀ receptor. The transfection system is also useful for determining the affinity and efficacy of known drugs at the human 5-HT₁₀

-23-

receptor sites.

This invention also provides a method of screening drugs to identify drugs which specifically interact with, and bind to, the human 5-HT_{1D} receptor on the surface of a cell which comprises contacting a mammalian cell comprising a DNA molecule encoding a human 5-HT_{1D} receptor on the surface of a cell with a plurality of drugs, determining those drugs which bind to the mammalian cell, and thereby identifying drugs which specifically interact with, and bind to, the human 5-HT_{1D} receptor. The DNA in the cell may have a coding sequence substantially the same as the coding sequence shown in Figure 3, or the coding sequence shown in Figure 4. Preferably, the mammalian cell is nonneuronal in origin. An example of a nonneuronal mammalian cell is an Ltk- cell, in particular the Ltk- cell designated L-5HT-8-30-84, or the Ltk- cell designated L-5HT-11. Drug candidates are identified by choosing chemical compounds which bind with high affinity to the expressed 5-HT_{1D} receptor protein in transfected cells, using radioligand binding methods well known in the art, examples of which are shown in the binding assays described herein. Drug candidates are also screened for selectivity by identifying compounds which bind with high affinity to one particular 5-HT_{1D} receptor subtype but do not bind with high affinity to any other serotonin receptor subtype or to any other known receptor site. Because selective, high affinity compounds interact primarily with the target 5-HT_{1D} receptor site after administration to the patient, the chances of producing a drug with unwanted side effects are minimized by this approach. This invention provides a pharmaceutical composition comprising a drug identified by the method described above and a pharmaceutically acceptable carrier.

-24-

Once the candidate drug has been shown to be adequately bio-available following a particular route of administration, for example orally or by injection (adequate therapeutic concentrations must be maintained at the site of action for an adequate period to gain the desired therapeutic benefit), and has been shown to be non-toxic and therapeutically effective in appropriate disease models, the drug may be administered to patients by that route of administration determined to make the drug bio-available, in an appropriate solid or solution formulation, to gain the desired therapeutic benefit.

This invention provides a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding a human 5-HT₁₀ receptor, for example with a coding sequence included within the sequence shown in Figure 3 or Figure 4. Nucleic acid probe technology is well known to those skilled in the art who will readily appreciate that such probes may vary greatly in length and may be labeled with a detectable label, such as a radioisotope or fluorescent dye, to facilitate detection of the probe. Detection of nucleic acid encoding human 5-HT₁₀ receptors is useful as a diagnostic test for any disease process in which levels of expression of the corresponding 5-HT₁₀ receptor is altered. DNA probe molecules are produced by insertion of a DNA molecule which encodes human 5-HT₁₀ receptor or fragments thereof into suitable vectors, such as plasmids or bacteriophages, followed by insertion into suitable bacterial host cells and replication and harvesting of the DNA probes, all using methods well known in the art. For example, the DNA may be extracted from a cell lysate using

35

-25-

phenol and ethanol, digested with restriction enzymes corresponding to the insertion sites of the DNA into the vector (discussed above), electrophoresed, and cut out of the resulting gel. Examples of such DNA molecules are shown in Figure 3 and in Figure 4. The probes are useful for 'in situ' hybridization or in order to locate tissues which express this gene family, or for other hybridization assays for the presence of these genes or their mRNA in various biological tissues. In addition, synthesized oligonucleotides (produced by a DNA synthesizer) complementary to the sequence of a DNA molecule which encodes human 5-HT_{1D} receptor or are useful as probes for these genes, for their associated mRNA, or for the isolation of related genes by homology screening of genomic or cDNA libraries, or by the use of amplification techniques such as the Polymerase Chain Reaction.

This invention also provides a method of detecting expression of a 5-HT_{1D} receptor on the surface of a cell by detecting the presence of mRNA coding for a 5-HT_{1D} receptor which comprises obtaining total mRNA from the cell using methods well known in the art and contacting the mRNA so obtained with a nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding a human 5-HT_{1D} receptor under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the 5-HT_{1D} receptor by the cell. Hybridization of probes to target nucleic acid molecules such as mRNA molecules employs techniques well known in the art. In one possible means of performing this method, nucleic acids are extracted by precipitation from lysed cells and the mRNA is

isolated from the extract using a column which binds the poly-A tails of the mRNA molecules. The mRNA is then exposed to radioactively labelled probe on a nitrocellulose membrane, and the probe hybridizes to and thereby labels complementary mRNA sequences. Binding may be detected by
5 autoradiography or scintillation counting. However, other methods for performing these steps are well known to those skilled in the art, and the discussion above is merely an example.

10 This invention provides an antisense oligonucleotide having a sequence capable of binding specifically with any sequences of an mRNA molecule which encodes a human 5-HT₁₀ receptor so as to prevent translation of the mRNA molecule. The antisense oligonucleotide may have a sequence capable of
15 binding specifically with any sequences of the cDNA molecule whose sequence is shown in Figure 3 or the cDNA molecule whose sequence is shown in Figure 4. A particular example of an antisense oligonucleotide is an antisense oligonucleotide comprising chemical analogues of
20 nucleotides.

This invention also provides a pharmaceutical composition comprising an amount of the oligonucleotide described above effective to reduce expression of a human 5-HT₁₀ receptor by
25 passing through a cell membrane and binding specifically with mRNA encoding a human 5-HT₁₀ receptor in the cell so as to prevent its translation and a pharmaceutically acceptable hydrophobic carrier capable of passing through a cell membrane. The oligonucleotide may be coupled to a substance
30 which inactivates mRNA, such as a ribozyme. The pharmaceutically acceptable hydrophobic carrier capable of passing through cell membranes may also comprise a structure

-27-

which binds to a receptor specific for a selected cell type and is thereby taken up by cells of the selected cell type. The structure may be part of a protein known to bind a cell-type specific receptor, for example an insulin molecule, which would target pancreatic cells. DNA molecules having coding sequences substantially the same as the coding sequence shown in Figure 3 or the coding sequence shown in Figure 4 may be used as the oligonucleotides of the pharmaceutical composition.

This invention also provides a method of treating abnormalities which are alleviated by reduction of expression of a 5-HT_{1D} receptor which comprises administering to a subject an amount of the pharmaceutical composition described above effective to reduce expression of the 5-HT_{1D} receptor by the subject. This invention further provides a method of treating an abnormal condition related to 5-HT_{1D} receptor activity which comprises administering to a subject an amount of the pharmaceutical composition described above effective to reduce expression of the 5-HT_{1D} receptor by the subject. Several examples of such abnormal conditions are dementia, Parkinson's disease, an eating disorder, pathological anxiety, or a migraine headache.

Antisense oligonucleotide drugs inhibit translation of mRNA encoding these receptors. Synthetic oligonucleotides, or other antisense chemical structures are designed to bind to mRNA encoding the 5-HT_{1D} receptor and inhibit translation of mRNA and are useful as drugs to inhibit expression of 5-HT_{1D} receptor genes in patients. This invention provides a means to therapeutically alter levels of expression of human 5-HT_{1D} receptors by the use of a synthetic antisense oligonucleotide drug (SAOD) which inhibits translation of

-28-

mRNA encoding these receptors. Synthetic oligonucleotides, or other antisense chemical structures designed to recognize and selectively bind to mRNA, are constructed to be complementary to portions of the nucleotide sequences shown in Figure 3 or Figure 4 of DNA, RNA or of chemically modified, artificial nucleic acids. The SAOD is designed to be stable in the blood stream for administration to patients by injection, or in laboratory cell culture conditions, for administration to cells removed from the patient. The SAOD is designed to be capable of passing through cell membranes in order to enter the cytoplasm of the cell by virtue of physical and chemical properties of the SAOD which render it capable of passing through cell membranes (e.g. by designing small, hydrophobic SAOD chemical structures) or by virtue of specific transport systems in the cell which recognize and transport the SAOD into the cell. In addition, the SAOD can be designed for administration only to certain selected cell populations by targeting the SAOD to be recognized by specific cellular uptake mechanisms which binds and takes up the SAOD only within certain selected cell populations. For example, the SAOD may be designed to bind to a receptor found only in a certain cell type, as discussed above. The SAOD is also designed to recognize and selectively bind to the target mRNA sequence, which may correspond to a sequence contained within the sequences shown in Figures 4 and 5, by virtue of complementary base pairing to the mRNA. Finally, the SAOD is designed to inactivate the target mRNA sequence by any of three mechanisms: 1) by binding to the target mRNA and thus inducing degradation of the mRNA by intrinsic cellular mechanisms such as RNase I digestion, 2) by inhibiting translation of the mRNA target by interfering with the binding of translation-regulating factors or of ribosomes, or 3) by inclusion of other chemical structures,

such as ribozyme sequences or reactive chemical groups, which either degrade or chemically modify the target mRNA. Synthetic antisense oligonucleotide drugs have been shown to be capable of the properties described above when directed against mRNA targets (J.S. Cohen, Trends in Pharm. Sci. 10, 435 (1989); H.M. Weintraub, Sci. Am. January (1990) p. 40). In addition, coupling of ribozymes to antisense oligonucleotides is a promising strategy for inactivating target mRNA (N. Sarver et al., Science 247, 1222 (1990)). An SAOD serves as an effective therapeutic agent if it is designed to be administered to a patient by injection, or if the patient's target cells are removed, treated with the SAOD in the laboratory, and replaced in the patient. In this manner, an SAOD serves as a therapy to reduce receptor expression in particular target cells of a patient, in any clinical condition which may benefit from reduced expression of 5-HT₁₀ receptors.

This invention provides an antibody directed to the human 5-HT₁₀ receptor, for example a monoclonal antibody directed to an epitope of a human 5-HT₁₀ receptor present on the surface of a cell and having an amino acid sequence substantially the same as an amino acid sequence for a cell surface epitope of the human 5-HT₁₀ receptor included in the amino acid sequence shown in Figure 3 or included in the amino acid sequence shown in Figure 4. Amino acid sequences may be analyzed by methods well known in the art to determine whether they produce hydrophobic or hydrophilic regions in the proteins which they build. In the case of cell membrane proteins, hydrophobic regions are well known to form the part of the protein that is inserted into the lipid bilayer which forms the cell membrane, while hydrophilic regions are located on the cell surface, in an aqueous environment.

-30-

Therefore antibodies to the hydrophilic amino acid sequences shown in Figure 3 or in Figure 4 will bind to a surface epitope of a human 5-HT₁₀ receptor, as described. Antibodies directed to human 5-HT₁₀ receptors may be serum-derived or monoclonal and are prepared using methods well known in the art. For example, monoclonal antibodies are prepared using hybridoma technology by fusing antibody producing B cells from immunized animals with myeloma cells and selecting the resulting hybridoma cell line producing the desired antibody. Cells such as SR3T3 cells or Ltk-cells may be used as immunogens to raise such an antibody. Alternatively, synthetic peptides may be prepared using commercially available machines and the amino acid sequences shown in Figure 3 and Figure 4. As a still further alternative, DNA, such as a cDNA or a fragment thereof, may be cloned and expressed and the resulting polypeptide recovered and used as an immunogen. These antibodies are useful to detect the presence of human 5-HT₁₀ receptors encoded by the isolated DNA, or to inhibit the function of the receptors in living animals, in humans, or in biological tissues or fluids isolated from animals or humans.

This invention provides a pharmaceutical composition which comprises an amount of an antibody directed to the human 5-HT₁₀ receptor effective to block binding of naturally occurring ligands to the 5-HT₁₀ receptor, and a pharmaceutically acceptable carrier. A monoclonal antibody directed to an epitope of a human 5-HT₁₀ receptor present on the surface of a cell and having an amino acid sequence substantially the same as an amino acid sequence for a cell surface epitope of the human 5-HT₁₀ receptor included in the amino acid sequence shown in Figure 3 or included in the amino acid sequence shown in Figure 4 are useful for this

-31-

purpose.

5 This invention also provides a method of treating abnormalities which are alleviated by reduction of expression of a human 5-HT₁₀ receptor which comprises
10 administering to a subject an amount of the pharmaceutical composition described above effective to block binding of naturally occurring ligands to the 5-HT₁₀ receptor and thereby alleviate abnormalities resulting from overexpression of a human 5-HT₁₀ receptor. Binding of the antibody to the
15 receptor prevents the receptor from functioning, thereby neutralizing the effects of overexpression. The monoclonal antibodies described above are both useful for this purpose. This invention additionally provides a method of treating an abnormal condition related to an excess of 5-HT₁₀ receptor
20 activity which comprises administering to a subject an amount of the pharmaceutical composition described above effective to block binding of naturally occurring ligands to the 5-HT₁₀ receptor and thereby alleviate the abnormal condition. Some examples of abnormal conditions are dementia, Parkinson's disease, an eating disorder, a pathological anxiety, and a migraine headache.

25 This invention provides a method of detecting the presence of a 5-HT₁₀ receptor on the surface of a cell which comprises contacting the cell with an antibody directed to the human 5-HT₁₀ receptor, under conditions permitting binding of the antibody to the receptor, detecting the presence of the antibody bound to the cell, and thereby the presence of the human 5-HT₁₀ receptor on the surface of the cell. Such a
30 method is useful for determining whether a given cell is defective in expression of 5-HT₁₀ receptors on the surface of the cell. Bound antibodies are detected by methods well

-32-

known in the art, for example by binding fluorescent markers to the antibodies and examining the cell sample under a fluorescence microscope to detect fluorescence on a cell indicative of antibody binding. The monoclonal antibodies described above are useful for this purpose.

5

This invention provides a transgenic nonhuman mammal expressing DNA encoding a human 5-HT₁₀ receptor. This invention also provides a transgenic nonhuman mammal expressing DNA encoding a human 5-HT₁₀ receptor so mutated as to be incapable of normal receptor activity, and not expressing native 5-HT₁₀ receptor. This invention also provides a transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a human 5-HT₁₀ receptor so placed as to be transcribed into antisense mRNA which is complementary to mRNA encoding a 5-HT₁₀ receptor and which hybridizes to mRNA encoding a 5-HT₁₀ receptor thereby reducing its translation. The DNA may additionally comprise an inducible promoter or additionally comprise tissue specific regulatory elements, so that expression can be induced, or restricted to specific cell types. Examples of DNA are DNA or cDNA molecules having a coding sequence substantially the same as the coding sequence shown in Figure 3, or the coding sequence shown in Figure 4. An example of a transgenic animal is a transgenic mouse. Examples of tissue specificity-determining regions are the metallothionein promoter (Low, M.J., Lechan, R.M., Hammer, R.E. et al. Science 231:1002-1004 (1986)) and the L7 promoter (Oberdick, J., Smeyne, R.J., Mann, J.R., Jackson, S. and Morgan, J.I. Science 248:223-226 (1990)).

30

Animal model systems which elucidate the physiological and behavioral roles of human 5-HT₁₀ receptors are produced by

35

-33-

creating transgenic animals in which the expression of a 5-HT₁₀ receptor is either increased or decreased, or the amino acid sequence of the expressed 5-HT₁₀ receptor protein is altered, by a variety of techniques. Examples of these techniques include: 1) Insertion of normal or mutant versions of DNA encoding a human 5-HT₁₀ receptor or homologous animal versions of these genes, by microinjection, retroviral infection or other means well known to those skilled in the art, into appropriate fertilized embryos in order to produce a transgenic animal (Hogan B. et al. Manipulating the Mouse Embryo, A Laboratory Manual, Cold Spring Harbor Laboratory (1986)). 2) Homologous recombination (Capecchi M.R. Science 244:1288-1292 (1989); Zimmer, A. and Gruss, P. Nature 338:150-153 (1989)) of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of expression or the structure of these 5-HT₁₀ receptors. The technique of homologous recombination is well known in the art. It replaces the native gene with the inserted gene and so is useful for producing an animal that cannot express native receptor but does express, for example, an inserted mutant receptor, which has replaced the native receptor in the animal's genome by recombination, resulting in underexpression of the receptor. Microinjection adds genes to the genome, but does not remove them, and so is useful for producing an animal which expresses its own and added receptors, resulting in overexpression of the receptor. One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as M2 medium (Hogan B. et al. Manipulating the Mouse Embryo, A Laboratory

-34-

Manual, Cold Spring Harbor Laboratory (1986)). DNA or cDNA encoding a human 5-HT₁₀ receptor is purified from a vector (such as plasmid p5HT-8-30-84 or p5HT-11 described above) by methods well known in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the trans-gene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the trans-gene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a pipet puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse (a mouse stimulated by the appropriate hormones to maintain pregnancy but which is not actually pregnant), where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg cell, and is used here only for exemplary purposes.

Since the normal action of receptor-specific drugs is to activate or to inhibit the receptor, the transgenic animal model systems described above are useful for testing the biological activity of drugs directed against these 5-HT₁₀ receptors even before such drugs become available. These animal model systems are useful for predicting or evaluating possible therapeutic applications of drugs which activate or inhibit these 5-HT₁₀ receptors by inducing or inhibiting expression of the native or trans-gene and thus increasing or decreasing expression of normal or mutant 5-HT₁₀ receptors in the living animal. Thus, a model system is produced in

-35-

which the biological activity of drugs directed against these 5-HT₁₀ receptors are evaluated before such drugs become available. The transgenic animals which over or under produce the 5-HT₁₀ receptor indicate by their physiological state whether over or under production of the 5-HT₁₀ receptor is therapeutically useful. It is therefore useful to evaluate drug action based on the transgenic model system. One use is based on the fact that it is well known in the art that a drug such as an antidepressant acts by blocking neurotransmitter uptake, and thereby increases the amount of neurotransmitter in the synaptic cleft. The physiological result of this action is to stimulate the production of less receptor by the affected cells, leading eventually to underexpression. Therefore, an animal which underexpresses receptor is useful as a test system to investigate whether the actions of such drugs which result in under expression are in fact therapeutic. Another use is that if overexpression is found to lead to abnormalities, then a drug which down-regulates or acts as an antagonist to 5-HT₁₀ receptor is indicated as worth developing, and if a promising therapeutic application is uncovered by these animal model systems, activation or inhibition of the 5-HT₁₀ receptor is achieved therapeutically either by producing agonist or antagonist drugs directed against these 5-HT₁₀ receptors or by any method which increases or decreases the expression of these 5-HT₁₀ receptors in man.

This invention provides a method of determining the physiological effects of expressing varying levels of human 5-HT₁₀ receptors which comprises producing a transgenic nonhuman animal whose levels of human 5-HT₁₀ receptor expression are varied by use of an inducible promoter which regulates human 5-HT₁₀ receptor expression. This invention

-36-

also provides a method of determining the physiological effects of expressing varying levels of human 5-HT₁₀ receptors which comprises producing a panel of transgenic nonhuman animals each expressing a different amount of human 5-HT₁₀ receptor. Such animals may be produced by introducing
5 different amounts of DNA encoding a human 5-HT₁₀ receptor into the oocytes from which the transgenic animals are developed.

This invention also provides a method for identifying a
10 substance capable of alleviating abnormalities resulting from overexpression of a human 5-HT₁₀ receptor comprising administering the substance to a transgenic nonhuman mammal expressing at least one artificially introduced DNA molecule encoding a human 5-HT₁₀ receptor and determining whether the
15 substance alleviates the physical and behavioral abnormalities displayed by the transgenic nonhuman mammal as a result of overexpression of a human 5-HT₁₀ receptor. Examples of DNA molecules are DNA or cDNA molecules having a coding sequence substantially the same as the coding
20 sequence shown in Figure 3, or the coding sequence shown in Figure 4.

This invention provides a pharmaceutical composition comprising an amount of the substance described supra
25 effective to alleviate the abnormalities resulting from overexpression of 5-HT₁₀ receptor and a pharmaceutically acceptable carrier.

This invention further provides a method for treating the
30 abnormalities resulting from overexpression of a human 5-HT₁₀ receptor which comprises administering to a subject an amount of the pharmaceutical composition described above

-37-

effective to alleviate the abnormalities resulting from overexpression of a human 5-HT₁₀ receptor.

5 This invention provides a method for identifying a substance capable of alleviating the abnormalities resulting from underexpression of a human 5-HT₁₀ receptor comprising administering the substance to the transgenic nonhuman mammal described above which expresses only nonfunctional human 5-HT₁₀ receptor and determining whether the substance
10 alleviates the physical and behavioral abnormalities displayed by the transgenic nonhuman mammal as a result of underexpression of a human 5-HT₁₀ receptor.

This invention also provides a pharmaceutical composition comprising an amount of a substance effective to alleviate
15 abnormalities resulting from underexpression of 5-HT₁₀ receptor and a pharmaceutically acceptable carrier.

This invention further provides a method for treating the abnormalities resulting from underexpression of a human 5-HT₁₀ receptor which comprises administering to a subject an
20 amount of the pharmaceutical composition described above effective to alleviate the abnormalities resulting from underexpression of a human 5-HT₁₀ receptor.

25 This invention provides a method for diagnosing a predisposition to a disorder associated with the expression of a specific human 5-HT₁₀ receptor allele which comprises:
a) obtaining DNA of subjects suffering from the disorder; b) performing a restriction digest of the DNA with a panel of
30 restriction enzymes; c. electrophoretically separating the resulting DNA fragments on a sizing gel; d) contacting the resulting gel with a nucleic acid probe capable of

-38-

specifically hybridizing to DNA encoding a human 5-HT₁₀ receptor and labelled with a detectable marker; e) detecting labelled bands which have hybridized to the the DNA encoding a human 5-HT₁₀ receptor labelled with a detectable marker to create a unique band pattern specific to the DNA of subjects suffering from the disorder; f) preparing DNA obtained for diagnosis by steps a-e; and g) comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step e and the DNA obtained for diagnosis from step f to determine whether the patterns are the same or different and thereby to diagnose predisposition to the disorder if the patterns are the same. This method may also be used to diagnose a disorder associated with the expression of a specific human 5-HT₁₀ receptor allele.

This invention provides a method of preparing the isolated 5-HT₁₀ receptor which comprises inducing cells to express 5-HT₁₀ receptor, recovering the receptor from the resulting cells, and purifying the receptor so recovered. An example of an isolated 5-HT₁₀ receptor is an isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 3. Another example of an isolated 5-HT₁₀ receptor is an isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 4. For example, cells can be induced to express receptors by exposure to substances such as hormones. The cells can then be homogenized and the receptor isolated from the homogenate using an affinity column comprising, for example, serotonin or another substance which is known to bind to the receptor. The resulting fractions can then be purified by contacting them with an ion exchange column, and determining which fraction contains receptor activity or binds anti-receptor

-39-

antibodies.

5 This invention provides a method of preparing the isolated 5-HT₁₀ receptor which comprises inserting nucleic acid encoding 5-HT₁₀ receptor in a suitable vector, inserting the
10 resulting vector in a suitable host cell, recovering the receptor produced by the resulting cell, and purifying the receptor so recovered. An example of an isolated 5-HT₁₀ receptor is an isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 3. Another example of an isolated 5-HT₁₀ receptor is
15 an isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 4. This method for preparing 5-HT₁₀ receptor uses recombinant DNA technology methods well known in the art. For example, isolated nucleic acid encoding 5-HT₁₀ receptor is inserted
20 in a suitable vector, such as an expression vector. A suitable host cell, such as a bacterial cell, or a eucaryotic cell such as a yeast cell, is transfected with the vector. 5-HT₁₀ receptor is isolated from the culture medium by affinity purification or by chromatography or by other methods well known in the art.

25 This invention provides an antisense oligonucleotide having a sequence capable of binding specifically with any sequences of an mRNA molecule which encodes a receptor so as to prevent translation of the mRNA molecule.

This invention also provides a transgenic nonhuman mammal expressing DNA encoding a receptor.

30 This invention further provides a transgenic nonhuman mammal expressing DNA encoding a receptor so mutated as to be

-40-

incapable of normal receptor activity, and not expressing native receptor.

5 This invention provides a method of determining the physiological effects of expressing varying levels of a receptor which comprises producing a transgenic nonhuman animal whose levels of receptor expression are varied by use of an inducible promoter which regulates receptor expression.

10 This invention also provides a method of determining the physiological effects of expressing varying levels of a receptor which comprises producing a panel of transgenic nonhuman animals each expressing a different amount of the receptor.

15 This invention further provides transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a receptor so placed as to be transcribed into antisense mRNA which is complementary to mRNA encoding the receptor and which hybridizes to mRNA encoding the receptor thereby preventing its translation.

20

25 This invention provides a method for determining whether a ligand not known to be capable of binding to a receptor can bind to a receptor which comprises contacting a mammalian cell comprising an isolated DNA molecule encoding the receptor with the ligand under conditions permitting binding of ligands known to bind to a receptor, detecting the presence of any of the ligand bound to the receptor, and thereby determining whether the ligand binds to the receptor.

30

-41-

Applicants have identified individual receptor subtype proteins and have described methods for the identification of pharmacological compounds for therapeutic treatments. Pharmacological compounds which are directed against specific receptor subtypes provide effective new therapies with minimal side effects.

The 5-HT₁₀ receptor subtype was first detected in the caudate nucleus of the bovine brain. It has been widely accepted that 5-HT₁₀ is the non-rodent analog of the 5-HT₁₈ receptor, which has a strong localization in basal ganglia (Schmidt and Peroutka, FASEB J. 3:2242 (1989)). Since the basal ganglia are involved in the control of movement, the 5-HT₁₀ receptor is important in movement control. Additionally, re-examination of previous reports of a 5-HT₁ receptor in Parkinson's disease patients, in light of the more recently described 5-HT₁₀ receptor pharmacology, yields an interesting observation. Parkinson's patients with dementia show a large decrease in 5-HT₁ receptors in the frontal cortex. (Cross, A.J., Crow, T.J., Johnston, J.A., Perry, E.K., Perry, R.H., Blessed, G. and Tomlinson, B.E., J. Neurol. Sci. 60: 383-392, (1989)). This region has recently been shown to contain mainly 5-HT₁₀ receptors within the 5-HT₁ class. (Herrick-Davis, K., Titeler, M., Leonhardt, S., Struble, R. and Price, D., J. Neurochem. 51: 1906-1912 (1988)). These data indicate that 5-HT₁₀ agonists will have significant therapeutic value in ameliorating the dementia of Parkinson's disease, and in treatment of other dementias involving 5-HT₁ receptors.

5-HT₁₀ receptors are implicated in the regulation of feeding behavior, thus providing a treatment for obesity and a potential insight into anorexia and bulimia. In rat models,

-42-

the 5-HT_{1D} receptor analog 5-HT_{1B} has been implicated in feeding. Agonists of the 5-HT_{1B} receptor such as RU 24924 (but not DPAT) and mCPP both decrease eating (Hutson, P.H., Kennett, G.A., Donohoe, T.P., Dourish, C.T., and Curzon, G. in Behavioral Pharmacology of 5-HT, Bevan, P., Cools, A.R., and Archer, T., eds. Lawrence Erlbaum Associates, Publishers; N.J. (1989) pp. 283-286; Samanin, R. in Behavioral Pharmacology of 5-HT, Bevan, P., Cools, A.R., and Archer, T., eds. Lawrence Erlbaum Associates, Publishers; N.J. (1989) pp. 259-283). In other studies, 5-HT_{1B} receptor antagonists cause anorexia (Leander, J.D., in Behavioral Pharmacology of 5-HT, Bevan, P., Cools, A.R., and Archer, T., eds. Lawrence Erlbaum Associates, Publishers; N.J. (1989) pp. 287-290). Localization of 5-HT_{1D} receptors in the nucleus accubens (Samanin, 1989; *ibid*), part of the neural substrate of feeding behavior, indicate that 5-HT_{1D} agonists have therapeutic value in the control of obesity. Furthermore, 5-HT_{1D} antagonists are useful in reversing endogenous anorexia and in the control of bulimic behavior.

5-HT_{1D} receptors are also implicated in anxiety. In the rat analog model, 5-HT_{1B} receptor agonists have been reported to decrease the startle response in rats and to effect the defensive burying behavior (Bevan, P., Lorens, S., and Archer, T. in Behavioral Pharmacology of 5-HT, Bevan, P., Cools, A.R., and Archer, T., eds. Lawrence Erlbaum Associates, Publishers; N.J. (1989) pp.459-474). Both paradigms indicate possible therapeutic roles of human 5-HT_{1D} agonists as anxiolytics. Furthermore, yohimbine (a 5-HT_{1D} antagonist) has been shown to worsen panic attacks in man (Charney, D.S., Henenger, G.R., and Breier, A., Arch. Gen. Psychiat. 41: 751-763, (1984)).

-43-

5 The best proven therapeutic application of 5-HT₁₀ receptor pharmacology is to migraine treatment. Sumatriptan, an agonist for certain 5-HT₁₀ receptor sites, has been demonstrated to be effective in the control of acute migraine attacks in clinical trials (Doenicke, A., Melchart, D.,
10 Bayliss, E.M., Cephalalgia 9 suppl 9:89-92 (1989)) with few side effects and good efficacy (Baer H.A., Brand, J., Doenicke, A., Melchart, D., Tryba, M., and Sahlender, H.M., Cephalalgia 9 (suppl 9): 83-87, (1989), Perrin, V.L., Farkkila, J., Goasguen, J., Donicke, A., Brand, J., and
15 Tfelt-Hansen, P., Cephalalgia 9 (suppl 9) 63-72, (1989)). Sumatriptan appears to be effective against not only the headache but also the nausea, vomiting and sensory disturbances suffered by migraine patients. Applicants have shown (Table 1) that sumatriptan has extremely high affinity
20 at the human 5-HT₁₀₋₁ and 5-HT₁₀₋₂ receptors.

25 This invention identifies for the first time two new receptor proteins, their amino acid sequences, and their human genes. Furthermore, this invention describes a previously unrecognized group of receptors within the definition of a 5-HT₁₀ receptor. The information and experimental tools provided by this discovery are useful to generate new therapeutic agents, and new therapeutic or
30 diagnostic assays for these new receptor proteins, their associated mRNA molecules or their associated genomic DNA. The information and experimental tools provided by this discovery will be useful to generate new therapeutic agents, and new therapeutic or diagnostic assays for these new receptor proteins, their associated mRNA molecules, or their associated genomic DNA.

Specifically, this invention relates to the first isolation

-44-

of human cDNA and genomic clones encoding 5-HT₁₀ receptors. Two new human genes for the receptors identified herein as 5-HT₁₀₋₁ and 5-HT₁₀₋₂ have been identified and characterized, and a series of related cDNA and genomic clones have been isolated. In addition, the human 5-HT₁₀ receptors have been expressed in Ltk- cells by transfecting the cells with the plasmids pcEXV-8-30-84 and pSVL-11. The pharmacological binding properties of the proteins encoded have been determined, and these binding properties classify these proteins as serotonin 5-HT₁₀ receptors. Mammalian cell lines expressing these human 5-HT₁₀ receptors at the cell surface have been constructed, thus establishing the first well-defined, cultured cell lines with which to study these 5-HT₁₀ receptors.

The invention will be better understood by reference to the Experimental Details which follow, but those skilled in the art will readily appreciate that the specific experiments detailed are only illustrative of the invention as described more fully in the claims which follow thereafter.

Experimental Details:Isolation of the dog RDC4 gene

The dog RDC4 gene (Libert et al. Science 244:569-572,1989) was isolated as a prerequisite to isolating clones for human 5-HT_{1D} genes. The RDC4 gene was obtained by isolating clones from a dog genomic library (Stratagene) with an oligonucleotide probe complementary to the large cytoplasmic loop of the RDC4 gene. Overlapping oligomers complementary to the dog RDC4 sequence (GenBank, accession number X14049) were labeled with ³²P-dATP and ³²P-dCTP by synthesis with the large fragment of DNA Polymerase (Maniatis et al. Molecular Cloning, Cold Spring Harbor, 1982). Clones showing positive hybridization to the probe were picked and inserts subcloned into the plasmid pUC-18 (Pharmacia, Piscataway, N.J.). Sequencing via the Sanger dideoxy method confirmed the isolation of a clone containing the entire coding region of the gene.

Isolation of two human 5-HT_{1D} genomic clones.

A human placental genomic library (Stratagene) was screened with the 1.3-kilobase (kb) Hind III-Sph I fragment from the dog RDC-4 clone. The probe was labeled with ³²P by the method of random priming (A.P. Feinberg and B. Vogelstein, Anal.Biochem. 137:266 (1984)). Hybridization was performed at 40°C in a solution containing 50% formamide, 10% dextran sulfate, 5X SSC (1X SSC is 0.15 M sodium chloride, 0.015 M sodium citrate), 1X Denhardt's (0.02% polyvinyl-pyrrolidone, 0.02% Ficoll, and 0.02% bovine serum albumin), and 200 µg/ml of sonicated salmon sperm DNA. The filters were washed at 50°C in 0.1X SSC containing 0.1% sodium dodecyl sulfate (SDS) and exposed at -70°C to Kodak XAR film in the presence of an intensifying screen. Lambda phage hybridizing to the

-46-

probe were plaque purified and DNA was prepared for Southern Blot analysis (Maniatis et al., Molecular Cloning, Cold Spring Harbor, 1982; E. Southern, J. Mol. Biol. 98:503, 1975). For subcloning and further Southern blot analysis, DNA was inserted into pUC18.

5

Isolation of cDNA clones

10

A human hippocampal cDNA library (Stratagene) was screened with an oligonucleotide probe derived from the dog RDC4 sequence (corresponding to amino acid positions number 165 to 188 of the RDC4 sequence in Figure 6, which falls within transmembrane region IV). Overlapping oligomers were labeled with ^{32}P -dATP and ^{32}P -dCTP as described above (Experimental Details) and the hybridizations and washes were performed as described above for genomic clones (Experimental Details).

15

DNA Sequencing

20

Nucleotide sequence analysis was done by the Sanger dideoxy nucleotide chain-termination method (S. Sanger, et al., Proc. Natl. Acad. Sci., 74: 5463-5467, 1977) on denatured double-stranded plasmid templates (Chen and Seeburg, DNA 4: 165, 1985) using Sequenase (U.S. Biochemical Corp., Cleveland, Ohio).

25

^3H -5HT Binding Assays

30

^3H -5HT (20.8-284 Ci/mMol; DuPont, NEN, Wilmington, DE) was used as a radioligand to detect the expression of the 5-HT gene products in membrane fractions isolated from either transiently or stably transfected cell lines (see Expression). The incubation buffer contained: 50mM Tris-Cl pH 7.4, 10mM MgSO_4 , 0.5 mM EDTA, 1% ascorbic acid and 0.1mM pargyline. Incubations were initiated by the addition of

35

-47-

cell membranes (10-50 μ g per well) to 96 well microtiter plates containing ^3H -5HT (final concentration 10^{-4} to 10^{-5}M) in a final volume of 250 μ l. After 15 minutes at 37°C (in the dark), the incubation was terminated by rapid filtration using a Brandel Cell Harvester (Model 48R; Brandel, Gaithersville, MD). Specific binding represented 70-80% of total binding, using 10^{-5}M 5-HT to define nonspecific binding. For competition studies, drugs were initially screened at a concentration of 1-10X their reported K_i values for 5-HT_{1D} receptor binding. Radioactivity trapped on GF/B filter strips was quantitated by liquid scintillation spectroscopy in a Beckman LS5000 TA scintillation counter using Ready Safe liquid scintillation cocktail (Beckman Instruments, Fullerton, CA) at an efficiency of 50-55%.

Experimental Results:

Isolation of genomic and cDNA clones encoding 5HT_{1D} receptors

We have screened a human genomic placental library with the 1.3 kb Hind III-Sph I restriction fragment derived from the dog RDC-4 clone. A total of five human clones were isolated and were characterized by restriction endonuclease mapping and DNA sequence analysis. By restriction analysis all five clones were judged to be different. By sequence analysis two clones, genomic clone hP-8 (human Placenta-8) and genomic clone hP-84, were highly homologous at the nucleotide level ($\approx 90\%$) to dog RDC4. Following sequence analysis, it was determined that genomic clone hP-8 contained sequence encoding the 5' untranslated (5'UT) region, the amino (NH₂) terminus, and transmembrane regions I-III (TM I-III) of a receptor which appears to represent the human homologue of dog RDC4. Genomic clone hP-84

-48-

5 contained sequence corresponding to TM V-VII as well as the
carboxy terminus and the 3' UT of this gene. Transmembrane
region IV was found to be present on a third clone, genomic
clone hP-30. These three genomic clones collectively span
the entire coding region of the gene and represent the human
version of RDC4. The gene for this new human receptor is
designated gene 8-30-84. cDNA clones corresponding to gene
8-30-84 have been isolated from a human hippocampus library
(Stratagene) and are designated cDNA clone hH-13 (human
Hippocampus-13) and cDNA clone hH-46 (Figure 1). cDNA
10 clones hH-13 and hH-46 collectively span the entire coding
region of this gene.

15 Two additional genomic clones, clone hP-10 and clone hP-11,
were also isolated and characterized. Although both clones
were homologous to clone RDC4 at the nucleotide level, their
homologies were not as strong ($\approx 75\%$ vs. $\approx 90\%$ for clones 8
and 84). While genomic clone hP-10 was found to represent
a pseudogene of the human RDC4 receptor gene (of gene 8-30-
84), genomic clone hP-11 exhibited a genomic structure which
20 appeared to be intronless in the coding region and encodes
a gene (gene 11) different from that of gene 8-30-84 (Figure
2).

Nucleotide sequence and deduced amino acid sequence of gene
25 8-30-84 (5-HT_{1D-1} gene) DNA sequence information obtained
from gene 8-30-84 is shown in Figure 3. An open reading
frame extending from an ATG start codon at position 1 to a
stop codon at position 1131 can encode a protein 376 amino
acids in length (peptide 8-30-84), having a relative
30 molecular mass (M_r) of 41,783. A comparison of this protein
sequence with previously characterized neurotransmitter
receptors indicates that gene 8-30-84 encodes a receptor

35

-49-

which is a new member of a family of molecules which span the lipid bilayer seven times and couple to guanine nucleotide regulatory proteins (the G protein-coupled receptor family). A variety of structural features which are invariant in this family were present in peptide 8-30-84. The greatest homology found was between peptide 8-30-84, the dog RDC-4 receptor and the 5-HT_{1A} receptor. A comparison of peptide 8-30-84 with the dog RDC4 sequence is shown in Figure 5, which displays a model of the receptor structure organized into seven transmembrane spanning domains. Overall, 87% sequence conservation between the dog RDC4 sequence and the human peptide 8-30-84 was observed over 378 amino acids. The greatest divergence between the dog RDC4 and human gene 8-30-84 protein sequences was seen at the extracellular amino terminus and the large cytoplasmic loop found between TM-V and TM-VI. The homology between these two receptors within the transmembrane regions alone was 92%, with TM-1 being the most divergent.

Nucleotide and deduced amino acid sequence of gene 11 (5-HT_{1D-2} gene)

DNA sequence information obtained from gene 11 is shown in Figure 4. An open reading frame extending from an ATG codon at position 1 to a stop codon at position 1198 can encode a protein 398 amino acids in length (peptide 11 having a relative molecular mass (M_r) of 44,333. A comparison of this protein sequence with previously characterized neurotransmitter receptors indicates that the protein product of gene 11 is also a new member of the G-protein-coupled receptor family. Homology profiles show that the receptor encoded by gene 11 (peptide 11) is tightly related to, but distinct from, the receptor encoded by gene 8-30-84 (peptide 8-30-84) and by the dog RDC4 gene, with the 5HT_{1A}

-50-

receptor being the next closest receptor with homology to peptide 11 (Figure 6). The overall amino acid homology between peptide 8-30-84 and peptide 11 is 56%, with a 76% amino acid homology within the transmembrane regions alone. Virtually no homology exists within the amino terminus, with slight homology found in the large cytoplasmic loop between TM-V and TM-VI as well as the carboxyl terminus. The most divergent of the transmembrane spanning domains is TM-1, which is only 54% identical between peptide 8-30-84 and peptide 11. There does exist a stretch of four amino acids within TM-1, however, which is identical between peptide 8-30-84 and peptide 11 as well as the dog RDC4 protein product. The sequence of these amino acids is serine(S), asparagine(N), alanine(A) and phenylalanine(F) and can be abbreviated as "SNAF". This sequence is unique to this subfamily of receptors because it diverges from a conserved sequence (GNxL) present in this position of TM-I in a variety of G-protein coupled receptors.

Receptor Expression in Transfected Mammalian Cells

In order to confirm the functional identity of the newly isolated gene we have expressed clones RDC4 and hP-11 in cultured cell lines. A DNA fragment containing the entire coding region of each clone was subcloned into the expression vector pSVL. The resulting plasmids, pcEXV-8-30-84 and pSVL-11 were transiently introduced into Cos-7 cells using the DEAE-dextran protocol (Cullen, Methods in Enz. 152: 684-704, 1987).

Stable cell lines were produced by cotransfection with the plasmid containing the bacterial gene aminoglycoside phosphotransferase into Ltk- cells (American Type Culture Collection, Rockville, MD, Cell Line CCL 1,3) using the

-51-

calcium phosphate technique (protocol & kit obtained from Specialty Media, Inc. Lavallette, NJ). Clones expressing aminoglycoside transferase were selected by the addition of 1 mg/ml G418 (Gibco Laboratories, Grand Island, NY) to the culture medium. ^3H -5HT was used to monitor 5-HT_{1D} receptor gene expression in these clones. Since ^3H -5HT can also bind to 5-HT_{1A}, 5-HT_{1B}, and 5-HT_{1C} receptors, the masking ligands pindolol (1 μM) and SCH 23390 (1 μM) were included in the incubation.

Cos-7 cells or Ltk- cells were pseudotransfected with vector not containing an insert in order to assess endogenous levels of ligand binding. At 2nM radioligand, no specific binding was detected. Therefore, Cos-7 and Ltk- cells provide useful models for transfection of a putative 5-HT_{1D} receptor. Transiently transfected Cos-7 cells bound ^3H -5-HT with high affinity (3 nM) and with an estimated site density of 0.63-1.28 pmole/mg protein for RDC4 and 0.602 to 0.95 pmole/mg protein for peptide 11. The presence of the masking ligands pindolol and SCH 23390 had no significant effect on the specific binding or the pharmacological profile of the detected binding sites in initial studies, and was therefore eliminated in subsequent experiments.

Ltk- cells transfected with RDC4, gene 8-30-84 or gene 11 bound ^3H -5-HT with high affinity (K_d = 3.6 nM, 4.0 nM or 2.3 nM, respectively). The estimated B_{max} is equal to 0.275 pm/mg protein for RDC4, 5.1 pmole/mg protein for gene 8-30-84 and 2.3 pmole/mg protein for gene 11. Further characterization was accomplished by performing competition experiments for a series of drugs. Analysis of the competition data was accomplished using the computer-assisted nonlinear regression program Accucomp (London

-52-

Software; Chagrin Falls, OH). Data are shown in Table 1 and indicate that binding detected in the RDC4, gene 8-30-84 or gene 11 transfected cells has properties expected for a 5-HT₁ receptor. Additionally, the differences in the affinity of a series of competitors for these two receptors indicates that the RDC4, as well as the human genes, gene 8-30-84 and gene 11, encode proteins with binding properties of a 5-HT₁₀ receptor, and thus uncovers the existence of at least two subtypes of the 5-HT₁₀ receptor.

10

15

20

25

30

35

53

TABLE 1: [³H]5-HT BINDING TO 5-HT_{1D} RECEPTORS

DRUG	Cloned Canine 5-HT _{1D-1} (RDC4)	Cloned Human 5-HT _{1D-1}	Cloned Human 5-HT _{1D-2}	Human Native Membranes 5-HT _{1D}
Lysergol	N.D.	0.63 ± 0.10	1.2 ± 0.1	---
5-Carboxamidotryptamine	0.92 ± 0.06	0.70 ± 0.06	1.6 ± 0.2	0.93 ^a
Methylergonovine	N.D.	0.86 ± 0.36	2.9 ± 1.5	---
Ergotamine	1.9 ± 0.3	1.0 ± 0.2	1.1 ± 0.2	0.3 ^b
CGS-12066B	N.D.	2.9 ± 0.3	2.1 ± 0.3	---
PAPP	35 ± 6	3.1 ± 0.2	10 ± 1	---
Sumatriptan	1.6 ± 0.2	3.4 ± 0.3	7.7 ± 0.5	---
Methysergide	7.7 ± 1.7	3.6 ± 0.6	25 ± 5	14 ^b
5-Hydroxytryptamine	5.0 ± 1.4	3.9 ± 0.3	4.3 ± 0.9	6.5 ^a
5-methoxy-DMT	5.1 ± 1.7	4.4 ± 0.6	21 ± 7	---

5-Methoxytryptamine	31 ± 11	4.8 ± 0.3	34 ± 6	6.2 ^a
Metergoline	N.D.	5.8 ± 1.1	18 ± 3	12 ^a
1-Naphthylpiperazine	N.D.	7.4 ± 0.6	12 ± 2	---
5-Benzoxyltryptamine	N.D.	9.6 ± 3.0	19 ± 8	---
Methiothepin	N.D.	11 ± 2	25 ± 7	17 ^c

55

DRUG	Cloned Canine 5-HT _{1D-1} (RDC4)	Cloned Human 5-HT _{1D-1}	Cloned Human 5-HT _{1D-2}	Human Brain Membranes 5-HT _{1D}
Dipropyl-5-CT	N.D.	13 ± 1	42 ± 4	---
Rauwolscine	5.3 ± 2.9	16 ± 2	40 ± 7	---
Yohimbine	32 ± 6	22 ± 2	27 ± 3	42 ^b
Ritanserlin	N.D.	54 ± 10	1200 ± 115	3600 ^c
TFMPP	112 ± 14	64 ± 5	114 ± 18	210 ^c
Ketanserlin	> 1000	73 ± 12	5198 ± 524	> 10000 ^b
Tryptamine	343 ± 15	86 ± 6	521 ± 82	46 ^c
DPAT	86 ± 21	120 ± 11	260 ± 32	340 ^b
NAN-190	N.D.	194 ± 12	652 ± 48	---
α-methyl-5-HT	N.D.	211 ± 82	133 ± 51	---
mCPP	329 ± 80	216 ± 20	361 ± 31	1200 ^c
SCH-23390	N.D.	676 ± 69	1214 ± 626	---

DOI	729 ± 102	714 ± 81	1667 ± 451	670 ^b
Phentolamine	N.D.	775 ± 72	2148 ± 685	---
2-methyl-5-HT	444 ± 30	915 ± 68	860 ± 109	1700 ^c
Spiperone	> 1000	995 ± 96	> 10000	> 10000 ^c
Pindolol	> 1000	4112 ± 504	4878 ± 302	> 10000 ^b
Corynanthine	N.D.	> 10000	> 10000	> 10000 ^c
Zacopride	> 10000	> 10000	> 10000	---

Estimated equilibrium dissociation constants (K_i values in nM) were calculated from the Cheng-Prusoff relationship for the indicated displacer compounds. Binding experiments were performed on cell membrane preparations from LM (tk-) cells stably transfected with the indicated clones.

^aWaeber, C., Schoeffter, P., Hoyer, D., and Palacios, J.M. Neurochem. Res., 15:567-582, 1990. ^bHerrick-Davis, K., Titeler, M., Leonhardt, S., Struble, R., and Price, D. J. Neurochem. 51:1906-1912, 1988.

^cPeroutka, S.J., Switzer, J.A., and Hamik, A. Synapse, 3:61-66, 1989. N.D., not determined.

-57-

Discussion

Applicants have cloned and characterized two DNA molecules encoding human 5-HT_{1D} receptors. The expression of these cDNA clones in Cos-7 cells and Ltk- cells results in the appearance of this type of receptor on the cell surface.

The starting point for this research was dog clone RDC4, which was originally reported by Libert et al. (Science 244:569, 1989) as one of a collection of G-protein coupled receptors cloned by PCR technology. Our analysis of the nucleotide sequence following the model of Strader, Sigal and Dixon (FASEB J. 3:1825 (1989)) indicated that RDC4 was likely to be a 5-HT receptor. In addition, RDC4 showed highest sequence homology to the human serotonin 5-HT_{1A} sequence (Libert et al., 1989). The RDC4 sequence was isolated from a dog genomic library in our laboratories, and was transfected into a mammalian cell line. No expression data on this clone have been reported previously by any laboratory.

RDC4-transfected Cos-7 and Ltk- cells were found to bind [³H]5-HT with apparent high affinity ($K_D \approx 3\text{nM}$; $B_{\text{max}} \approx 0.25\text{--}0.4\text{pm/mg protein}$). This property indicates that RDC4 encodes a 5-HT₁ receptor. High affinity ³H5-HT binding was not affected by co-incubation with (-) pindolol (1 μM). This rules out the possibility that binding is to a 5-HT_{1A} or 5-HT_{1B} receptor. Furthermore, the 5-HT_{1C} antagonist SCH 23390 (1 μM) is also ineffective in reducing the binding of ³H-5-HT. Other compounds unable to compete for ³H-5HT binding sites are ketanserin (5-HT₂ receptors), ICS 205-930 (5-HT₃ and 5-HT₄), and 6-OH-indalpine (5-HT_{1P}). Furthermore, the

-58-

high affinity of 5-CT for this binding site rules out the possibility that binding is to a 5-HT_{1E} receptor. The strong sequence homology between RDC4 and the cloned human gene 8-30-84 indicate that the human gene, when transfected into a mammalian cell line, should express very similar pharmacological properties. This has been verified (see Table 1).

RDC4 was also used to clone a second human gene, gene 11. This gene was sequenced and expressed transiently in Cos-7 cells and as a stable cell line in Ltk-. As was shown to be true for the RDC4 clone, gene 11- transfected cells were found to bind ³H-5-HT with high affinity, and to be insensitive to compounds having affinity for 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1C}, 5-HT₂, 5-HT₃, or 5-HT₄ receptors. Additionally, the ability of low concentrations of 5-CT to compete with [³H]5-HT for binding to gene 11-transfected cells excludes the possibility that this gene encodes a 5-HT_{1P} or 5-HT_{1E} receptor. The pharmacological profile is shown in Table 1. These data are consistent with the characterization of gene 11 as encoding a 5-HT_{1D} receptor.

Table 1 demonstrates that although RDC4 (and peptide 8-30-84) and peptide 11 can both be classified as 5-HT_{1D} receptors, they have distinct pharmacological differences as well as distinct structures. These data demonstrate for the first time that the human 5-HT_{1D} binding site constitutes not a single receptor but a family of receptors. Potential differences in localization as well as pharmacology may be exploited in the future to develop powerful new drugs for selectively activating or inhibiting each of these 5-HT_{1D} receptor sites.

-59-

In summary, we have cloned two human 5-HT_{1D} receptors with differing pharmacological properties. This indicates that 5-HT_{1D} receptors constitute a receptor family rather than a single protein. In human brain, 5-HT_{1D} receptors have been shown to have the strongest representation in frontal cortex, putamen, caudate, both parts of the globus pallidus and the substantia nigra. Lower amounts are found in the raphe nuclei, hippocampus, and accumbens. 5-HT_{1D} receptors have been implicated in a number of clinically important conditions including migraine, appetite, movement control, anxiety, and dementia. 5-HT_{1D} agonists such as sumatriptan, are now in clinical trials. Our data, showing that there are multiple subtypes of human 5-HT_{1D} receptors, opens a new avenue for drug discovery. Pharmacological differences in these 5-HT_{1D} receptor subtypes, coupled with our exploration of possible differences in their localization, will likely lead to the development of more selective therapeutic agents with differential sensitivity for these subtypes.

Additional Experimental Data Relating to Pharmacological Characterization of a Cloned Rat 5-HT_{1B} Receptor: Relationship to the Cloned 5-HT_{1D} Receptor

The 5-HT_{1B} and 5-HT_{1D} receptors display similarities in their pharmacology, second messenger coupling, and anatomical distribution. This close relationship has led to the proposal that these receptors are species homologs, whereby the rat and mouse (and opossum) express the 5-HT_{1B} receptor and other species such as human, dog, bovine and guinea pig express the 5-HT_{1D} receptor (Hoyer and Middlemiss, 1989). The recent cloning of a 5-HT_{1D} receptor (Branchek et al., 1990) has allowed access to a direct assessment of the relationship between these two receptors using molecular cloning techniques. To isolate the putative rat homolog of

-60-

the 5-HT_{1D} gene, a rat library was screened at high stringency using a human 5-HT_{1D} sequence as a probe. A strongly hybridizing signal was selected for study, subcloned, and transiently transfected into a mammalian cell line (Cos-7). To identify the pharmacological profile of this receptor, radioligand binding assays were performed on a membrane fraction derived from the transfected cells. [³H]5-HT and [¹²⁵I]iodocyanopindolol ([¹²⁵I]I-CYP in the presence of 3 μM isoproterenol to block endogenous β-adrenergic receptors) were used as radioligands. [¹²⁵I]iodocyanopindolol was found to bind to membranes derived from the transfected cells with subnanomolar affinity [$K_d = 0.16$ nM]. In contrast, the equilibrium dissociation constant for [³H]5-HT binding was 18.4 nM. These values were in marked contrast to those measured using the human 5-HT_{1D} clone ([³H]5-HT: $K_d = 4$ nM; [¹²⁵I]I-CYP: no specific binding detected at 1 nM). The rauwolfia alkaloid rauwolscine was essentially inactive ($K_i > 10,000$ nM) at the rat receptor as was dipropyl-5-carboxyamidotryptamine. The rank order of potencies for the compounds studied was 5-HT > (-) propranolol > 5 methoxytryptamine > tryptamine > dipropyl-5-carboxyamidotryptamine = rauwolscine. These data clearly indicate that the gene isolated by homology with the human 5-HT_{1D} receptor encodes a protein with pharmacological properties which identify it as a 5-HT_{1B} receptor. Therefore, it appears that these receptors are species homologs which exhibit significant differences in pharmacology. A similar situation has been described for the rat and human 5-HT₂ receptors which have been cloned previously and shown to be highly homologous (Hartig et al., 1990). However, the differences in the pharmacological properties of the 5-HT_{1B}/5-HT_{1D} receptors have been shown to

-61-

be more robust than those of the rat 5-HT₂/human 5-HT₂ receptors.

5

10

15

20

25

30

35

-62-

What is claimed is:

1. An isolated nucleic acid molecule encoding a human 5-HT_{1D} receptor.
- 5 2. An isolated DNA molecule of claim 1.
3. A cDNA molecule of claim 2 encoding a human 5-HT_{1D-1} receptor having a coding sequence substantially the same as the coding sequence shown in Figure 3.
- 10 4. A cDNA molecule of claim 2 encoding a human 5-HT_{1D-2} receptor having a coding sequence substantially the same as the coding sequence shown in Figure 4.
- 15 5. An isolated protein which is a human 5-HT_{1D} receptor.
6. An isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 3.
- 20 7. An isolated protein having substantially the same amino acid sequence as the amino acid sequence shown in Figure 4.
- 25 8. A vector comprising the DNA molecule of claim 2.
9. A plasmid comprising the vector of claim 8.
- 30 10. A vector of claim 8 adapted for expression in a bacterial cell which comprises the regulatory elements necessary for expression of the DNA in the bacterial cell so located relative to the DNA encoding the 5-HT_{1D}

-63-

receptor as to permit expression thereof.

5 11. A vector of claim 8 adapted for expression in a yeast cell which comprises the regulatory elements necessary for expression of the DNA in the yeast cell so located relative to the DNA encoding the 5-HT_{1D} receptor as to permit expression thereof.

10 12. A vector of claim 8 adapted for expression in a mammalian cell which comprises the regulatory elements necessary for expression of the DNA in the mammalian cell so located relative to the DNA encoding the 5-HT_{1D} receptor as to permit expression thereof.

15 13. A plasmid of claim 9 adapted for expression in a mammalian cell which comprises the regulatory elements necessary for expression of the DNA in the mammalian cell so located relative to the DNA encoding the 5-HT_{1D} receptor as to permit expression thereof.

20 14. A plasmid comprising the cDNA molecule of claim 3 adapted for expression in a mammalian cell which comprises the regulatory elements necessary for expression of the DNA in the mammalian cell so located relative to the cDNA molecule as to permit expression thereof, designated pcEXV-8-30-84 (ATCC Accession No. 40790).

25

30 15. A plasmid comprising the cDNA molecule of claim 4 adapted for expression in a mammalian cell which comprises the the regulatory elements necessary for expression of the DNA in the mammalian cell so located relative to the cDNA molecule as to permit expression

-64-

thereof, designated pSVL-11 (ATCC Accession No. 40791).

16. A mammalian cell comprising the plasmid of claim 9.
17. An Ltk- cell comprising the plasmid of claim 9.
- 5 18. An Ltk- cell comprising the plasmid of claim 14, designated Ltk-8-30-84 (ATCC Accession No. CRL 10421).
- 10 19. An Ltk- cell comprising the plasmid of claim 15, designated Ltk-11 (ATCC Accession No. CRL 10422).
- 15 20. A method for determining whether a ligand not known to be capable of binding to a human 5-HT₁₀ receptor can bind to a human 5-HT₁₀ receptor which comprises contacting a mammalian cell comprising an isolated DNA molecule encoding a human 5-HT₁₀ receptor with the ligand under conditions permitting binding of ligands known to bind to a 5-HT₁₀ receptor, detecting the presence of any of the ligand bound to a human 5-HT₁₀ receptor, and thereby determining whether the ligand binds to a human 5-HT₁₀ receptor.
- 20 21. The method of claim 20 wherein the mammalian cell is nonneuronal in origin.
- 25 22. A method of claim 21, wherein the mammalian cell nonneuronal in origin is an Ltk- cell.
23. A ligand detected by the method of claim 20.
- 30 24. A method of screening drugs to identify drugs which specifically interact with, and bind to, the human 5-

-65-

HT₁₀ receptor on the surface of a cell which comprises contacting a mammalian cell comprising an isolated DNA molecule encoding a human 5-HT₁₀ receptor with a plurality of drugs, determining those drugs which bind to the mammalian cell, and thereby identifying drugs which specifically interact with, and bind to, a human 5-HT₁₀ receptor.

25. The method of claim 24 wherein the mammalian cell is nonneuronal in origin.

26. The method of claim 25 wherein the mammalian cell nonneuronal in origin is an Ltk- cell.

27. A pharmaceutical composition comprising a drug identified by the method of claim 24 and a pharmaceutically acceptable carrier.

28. A nucleic acid probe comprising a nucleic acid molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence of a nucleic acid molecule encoding a human 5-HT₁₀ receptor.

29. A DNA probe comprising a DNA molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence shown in Figure 3.

30. A DNA probe comprising a DNA molecule of at least 15 nucleotides capable of specifically hybridizing with a sequence included within the sequence shown in Figure 4.

-66-

- 5 31. A method of detecting expression of the 5-HT_{1D} receptor on the surface of a cell by detecting the presence of mRNA coding for a 5-HT_{1D} receptor which comprises obtaining total mRNA from the cell and contacting the mRNA so obtained with the nucleic acid probe of claim 28 under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the 5-HT_{1D} receptor by the cell.
- 10 32. An antisense oligonucleotide having a sequence capable of binding specifically with any sequences of an mRNA molecule which encodes a human 5-HT_{1D} receptor so as to prevent translation of the mRNA molecule.
- 15 33. An antisense oligonucleotide having a sequence capable of binding specifically with any sequences of the cDNA molecule of claim 3.
- 20 34. An antisense oligonucleotide having a sequence capable of binding specifically with any sequences of the cDNA molecule of claim 4.
- 25 35. An antisense oligonucleotide of claim 32 comprising chemical analogues of nucleotides.
- 30 36. A pharmaceutical composition comprising an amount of the oligonucleotide of claim 32 effective to reduce expression of a human 5-HT_{1D} receptor by passing through a cell membrane and binding specifically with mRNA encoding a human 5-HT_{1D} receptor in the cell so as to prevent its translation and a pharmaceutically acceptable hydrophobic carrier capable of passing through a cell membrane.

-67-

37. A pharmaceutical composition of claim 36, wherein the oligonucleotide is coupled to a substance which inactivates mRNA.
- 5 38. A pharmaceutical composition of claim 37, wherein the substance which inactivates mRNA is a ribozyme.
- 10 39. A pharmaceutical composition of claim 36, wherein the pharmaceutically acceptable hydrophobic carrier capable of passing through a cell membrane comprises a structure which binds to a receptor specific for a selected cell type and is thereby taken up by cells of the selected cell type.
- 15 40. A method of treating abnormalities which are alleviated by reduction of expression of a 5-HT₁₀ receptor which comprises administering to a subject an amount of the pharmaceutical composition of claim 36 effective to reduce expression of the 5-HT₁₀ receptor by the subject.
- 20 41. A method of treating an abnormal condition related to an excess of 5-HT₁₀ receptor activity which comprises administering to a subject an amount of the pharmaceutical composition of claim 36 effective to reduce expression of the 5-HT₁₀ receptor by the subject.
- 25 42. The method of claim 41 wherein the abnormal condition is dementia.
- 30 43. The method of claim 41 wherein the abnormal condition is Parkinson's disease.

-68-

44. The method of claim 41 wherein the abnormal condition is an eating disorder.
45. The method of claim 41 wherein the abnormal condition is a pathological anxiety.
- 5 46. The method of claim 41 wherein the abnormal condition is a migraine headache.
47. An antibody directed to a human 5-HT₁₀ receptor.
- 10 48. A monoclonal antibody directed to an epitope of a human 5-HT₁₀ receptor present on the surface of a cell and having an amino acid sequence substantially the same as an amino acid sequence for a cell surface epitope of the human 5-HT₁₀₋₁ receptor included in the amino acid sequence shown in Figure 3.
- 15 49. A monoclonal antibody directed to an epitope of a human 5-HT₁₀ receptor present on the surface of a cell and having an amino acid sequence substantially the same as an amino acid sequence for a cell surface epitope of the human 5-HT₁₀₋₂ receptor included in the amino acid sequence shown in Figure 4.
- 20 50. A pharmaceutical composition which comprises an amount of the antibody of claim 47 effective to block binding of naturally occurring ligands to the 5-HT₁₀ receptor and a pharmaceutically acceptable carrier.
- 25 51. A method of treating abnormalities which are alleviated by reduction of expression of a human 5-HT₁₀ receptor which comprises administering to a subject an amount of
- 30

-69-

the pharmaceutical composition of claim 50 effective to block binding of naturally occurring ligands to the 5-HT_{1D} receptor and thereby alleviate abnormalities resulting from overexpression of a human 5-HT_{1D} receptor.

5

52. A method of treating an abnormal condition related to an excess of 5-HT_{1D} receptor activity which comprises administering to a subject an amount of the pharmaceutical composition of claim 50 effective to
10 block binding of naturally occurring ligands to the 5-HT_{1D} receptor and thereby alleviate the abnormal condition.

53. The method of claim 52 wherein the abnormal condition
15 is dementia.

54. The method of claim 52 wherein the abnormal condition is Parkinson's disease.

20 55. The method of claim 52 wherein the abnormal condition is an eating disorder.

56. The method of claim 52 wherein the abnormal condition is a pathological anxiety.

25

57. The method of claim 52 wherein the abnormal condition is a migraine headache.

30

58. A method of detecting the presence of a human 5-HT_{1D} receptor on the surface of a cell which comprises contacting the cell with the antibody of claim 47 under conditions permitting binding of the antibody to the

-70-

receptor, detecting the presence of the antibody bound to the cell, and thereby detecting the presence of a human 5-HT_{1D} receptor on the surface of the cell.

- 5 59. A transgenic nonhuman mammal expressing DNA encoding a human 5-HT_{1D} receptor.
- 10 60. A transgenic nonhuman mammal expressing DNA encoding a human 5-HT_{1D} receptor so mutated as to be incapable of normal receptor activity, and not expressing native 5-HT_{1D} receptor.
- 15 61. A transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a human 5-HT_{1D} receptor so placed as to be transcribed into antisense mRNA which is complementary to mRNA encoding a 5-HT_{1D} receptor and which hybridizes to mRNA encoding a 5-HT_{1D} receptor thereby reducing its translation.
- 20 62. The transgenic nonhuman mammal of any of claims 59, 60 or 61, wherein the DNA encoding a human 5-HT_{1D} receptor additionally comprises an inducible promoter.
- 25 63. The transgenic nonhuman mammal of any of claims 59, 60 or 61, wherein the DNA encoding a human 5-HT_{1D} receptor additionally comprises tissue specific regulatory elements.
- 30 64. A transgenic mouse of any of claims 59, 60 or 61.
65. A method of determining the physiological effects of expressing varying levels of human 5-HT_{1D} receptors which comprises producing a transgenic nonhuman animal

-71-

whose levels of human 5-HT₁₀ receptor expression are varied by use of an inducible promoter which regulates human 5-HT₁₀ receptor expression.

5 66. A method of determining the physiological effects of expressing varying levels of human 5-HT₁₀ receptors which comprises producing a panel of transgenic nonhuman animals each expressing a different amount of human 5-HT₁₀ receptor.

10 67. A method for identifying a substance capable of alleviating the abnormalities resulting from overexpression of a human 5-HT₁₀ receptor comprising administering a substance to the transgenic nonhuman mammal of claim 59 and determining whether the
15 substance alleviates the physical and behavioral abnormalities displayed by the transgenic nonhuman mammal as a result of overexpression of a human 5-HT₁₀ receptor.

20 68. A pharmaceutical composition comprising an amount of a substance effective to alleviate the abnormalities resulting from overexpression of a human 5-HT₁₀ receptor and a pharmaceutically acceptable carrier.

25 69. A method for treating the abnormalities resulting from overexpression of a human 5-HT₁₀ receptor which comprises administering to a subject an amount of the pharmaceutical composition of claim 68 effective to
30 alleviate the abnormalities resulting from overexpression of a human 5-HT₁₀ receptor.

70. A method for identifying a substance capable of

-72-

alleviating the abnormalities resulting from underexpression of a human 5-HT_{1D} receptor comprising administering the substance to the transgenic nonhuman mammal of either of claims 60 or 61 and determining whether the substance alleviates the physical and behavioral abnormalities displayed by the transgenic nonhuman mammal as a result of underexpression of a human 5-HT_{1D} receptor.

71. A pharmaceutical composition comprising an amount of a substance effective to alleviate abnormalities resulting from underexpression of 5-HT_{1D} receptor and a pharmaceutically acceptable carrier.
72. A method for treating the abnormalities resulting from underexpression of a human 5-HT_{1D} receptor which comprises administering to a subject an amount of the pharmaceutical composition of claim 71 effective to alleviate the abnormalities resulting from underexpression of a human 5-HT_{1D} receptor.
73. A method for diagnosing a predisposition to a disorder associated with the expression of a specific human 5-HT_{1D} receptor allele which comprises:
- obtaining DNA of subjects suffering from the disorder;
 - performing a restriction digest of the DNA with a panel of restriction enzymes;
 - electrophoretically separating the resulting DNA fragments on a sizing gel;

-73-

- d. contacting the resulting gel with a nucleic acid probe capable of specifically hybridizing to DNA encoding a human 5-HT₁₀ receptor and labelled with a detectable marker;
- 5 e. detecting labelled bands which have hybridized to the the DNA encoding a human 5-HT₁₀ receptor labelled with a detectable marker to create a unique band pattern specific to the DNA of subjects suffering from the disorder;
- 10 f. preparing DNA obtained for diagnosis by steps a-e; and
- 15 g. comparing the unique band pattern specific to the DNA of subjects suffering from the disorder from step e and the DNA obtained for diagnosis from step f to determine whether the patterns are the same or different and to diagnose thereby predisposition to the disorder if the patterns are the same.
- 20
74. The method of claim 73 wherein a disorder associated with the expression of a specific human 5-HT₁₀ receptor allele is diagnosed.
- 25
75. A method of preparing the isolated 5-HT₁₀ receptor of claim 5 which comprises:
- 30 a. inducing cells to express 5-HT₁₀ receptor;
- b. recovering the receptor from the resulting cells; and

c. purifying the receptor so recovered.

76. A method of preparing the isolated 5-HT_{1D} receptor of claim 5 which comprises:

5 a. inserting nucleic acid encoding 5-HT_{1D} receptor in a suitable vector;

b. inserting the resulting vector in a suitable host cell;

10 c. recovering the receptor produced by the resulting cell; and

d. purifying the receptor so recovered.

15 77. An antisense oligonucleotide having a sequence capable of binding specifically with any sequences of an mRNA molecule which encodes a receptor so as to prevent translation of the mRNA molecule.

20 78. A transgenic nonhuman mammal expressing DNA encoding a receptor.

25 79. A transgenic nonhuman mammal expressing DNA encoding a receptor so mutated as to be incapable of normal receptor activity, and not expressing native receptor.

30 80. A method of determining the physiological effects of expressing varying levels of a receptor which comprises producing a transgenic nonhuman animal whose levels of receptor expression are varied by use of an inducible promoter which regulates receptor expression.

-75-

81. A method of determining the physiological effects of expressing varying levels of a receptor which comprises producing a panel of transgenic nonhuman animals each expressing a different amount of 5-HT_{1D} receptor.
- 5 82. A transgenic nonhuman mammal whose genome comprises antisense DNA complementary to DNA encoding a receptor so placed as to be transcribed into antisense mRNA which is complementary to mRNA encoding the receptor and which hybridizes to mRNA encoding the receptor
10 thereby preventing its translation.
83. A method for determining whether a ligand not known to be capable of binding to a receptor can bind to a receptor which comprises contacting a mammalian cell
15 comprising an isolated DNA molecule encoding the receptor with the ligand under conditions permitting binding of ligands known to bind to a receptor, detecting the presence of any of the ligand bound to the receptor, and thereby determining whether the
20 ligand binds to the receptor.


25

30

35


1/10

FIGURE 1



2/10

FIGURE 2

Gene II - A Related Gene

3 / 10.

-280 -260 -240
TGTGGTTTAGGAAAGACCTTTAACTACCAGCTGGTAGTTGTCTCAGCATTCTTCAAATAG

-220 -200 -180
TCCGGTCTTGTTTAATAATATTATTATTATTGTTATTTAATTTTATTGCAACTGT

-160 -140 -120
ACTTAGAGAATAGTCTGGTCTTGAGACCTTTTCACTGTGGTCTGTTCTGGTGTACGGCTC

-100 -80 -60
CCACCAGTGTGAAGCAGAAGGATGACTTTGCTCTGTTGTCTCAGGACAACCTTGAAGGAAGG

-40 -20 0
AGCCAAATGTGTGGAGGTCTGTGGGAAGAGAGAGCCACCTAGCATGTCCCCACTGAACCA
M S P L N Q

20 40 60
GTCAGCAGAAGGCCTTCCCCAGGAGGCCTCCAACAGATCCCTGAATGCCACAGAAACCTC
S A E G L P Q E A S N R S L N A T E T S

80 100 120
AGAGGCTTGGGATCCCAGGACCCTCCAGGCTCTCAAGATCTCCCTTCCCGTGCTCCTTTC
E A W D P R T L Q A L K I S L P V L L S

140 160 180
CGTCATCACACTGGCCACAGTCCTCTCCAATGCCTTTGTACTCACCACCATCTTACTCAC
V I T L A T V L S N A F V L T T I L L T

200 220 240
CAGGAAGCTCCACACCCCTGCCAACTACCTGATTGGCTCCCTGGCCACCACCGACCTCTT
R K L H T P A N Y L I G S L A T T D L L

260 280 300
GGTTTCCATCTTGGAATGCCCATCAGCATGGCCTATACCATCACCCACACCTGGAACCTT
V S I L V M P I S M A Y T I T H T W N F

320 340 360
TGGCCAAATCTTGTGTGACATCTGGCTGTCTCTGACATCACGTGCTGCACAGCCTCCAT
G Q I L C D I W L S S D I T C C T A S I

380 400 420
CCTGCATCTCTGTGTCATTGCTCTGGACAGGTACTGGGCAATCACAGATGCCCTGGAATA
L H L C V I A L D R Y W A I T D A L E Y

440 460 480
CAGTAAACGCAGGACGGCTGGCCACGCGGCCACCATGATCGCCATTGTCTGGGCCATCTC
S K R R T A G H A A T M I A I V W A I S

500 520 540
CATCTGCATCTCCATCCCCCGCTCTTCTGGCGGCAGGAAAAGGCCAGGAGGAGATGTC
I C I S I P P L F W R Q E K A Q E E M S

560 580 600
GGACTGTCTGGTGAACACCTCTCAGATCTCCTACACCATCTACTCCACCTGTGGGGCCTT
D C L V N T S O I S Y T I Y S T C G A F

Figure 3 (cont'd)

620 640 660
CTACATTCCCTCGGTGTTGCTCATCATCCTATATGGCCGGATCTACCGGGCTGCCCGGAA
Y I P S V L L I I L Y G R I Y R A A R N

680 700 720
CCGCATCCTGAATCCACCCTCACTCTCTGTTGGAAGCGCTTACCACGGCCACCTCATCAC
R I L N P P S L S G K R F T T A H L I T

740 760 780
AGGCTCTGCCGGGTCCGTCTGCTCGCTCAACTCCAGCCTCCATGAGGGGCACTCGCACTC
G S A G S V C S L N S S L H E G H S H S

800 820 840
GGCTGGCTCCCCTCTCTTTTCAACCACGTGAAAATCAAGCTTGCTGACAGTGCCCTGGA
A G S P L F F N H V K I K L A D S A L E

860 880 900
ACGCAAGAGGATTTCTGCTGCTCGAGAAAGGAAAGCCACTAAAATCCTGGGCATCATTCT
R K R I S A A R E R K A T K I L G I I L

920 940 960
GGGGGCCTTTATCATCTGCTGGCTGCCCTTCTTCGTGGTGTCTCTGGTCCTCCCCATCTG
G A F I I C W L P F F V V S L V L P I C

980 1000 1020
CCGGGACTCCTGCTGGATCCACCCGGGGCTCTTTGACTTCTTCACCTGGCTAGGCTATTT
R D S C W I H P G L F D F F T W L G Y L

1040 1060 1080
AAACTCCCTCATCAATCCAATAATCTACACTGTGTTTAATGAAGAGTTTCGGCAAGCTTT
N S L I N P I I Y T V F N E E F R Q A F

1100 1120 1140
TCAGAAAATTGTCCCTTTCGGAAGGCCTCCTAGTCTTATTCGATGAGGTAAAGAAACCT
Q K I V P F R K A S

1160 1180 1200
GCTTATGGGCTGGGCATGGTGGCTCATGCCTGTGATCCCAGCACTTTGGGAAGCTGAAGA

1220 1240 1260
GGAGGACTACTTGAGCTCAGGAGTTTGAGACCAGCCTGGGCAGCATAGGGAGACCCCGTC

1280 1300 1320
TCTAACGACAACAACAGAAATTACCGGACATGGTGGAGCGCGCCAGTAGTCCCAGCTACT

1340
CGAGAGGGCTGTGGCTACAGTGA

Figure 4

-110 -90 -70
AATGAGGAACACGGAACACAACGGGAACAAACCGCGACGCCACGGTCCTTCCGCCCTCT
-50 -30 -10
CCTTCGTCCGCTCCATGCCCAAGAGCTGAGCTCCGGAGCTGGGGCGAGGAGAGCCATGGA
M E
10 30 50
GGAACCGGGTGCTCAGTGCCTCCACCGGCGCCAGCGGGCTCCGAGACCTGGGTTCTCA
E P G A Q C A P P A P A G S E T W V P Q
70 90 110
AGCCAACTTATCCTCTGCTCCCTCCCAAACCTGCAGCGCCAAGGACTACATTTACCAGGA
A N L S S A P S Q N C S A K D Y I Y Q D
130 150 170
CTCCATCTCCCTACCCTGGAAAGTACTGCTGGTTATGCTATTGGCGCTCATCACCTTGGC
S I S L P W K V L L V M L L A L I T L A
190 210 230
CACCACGCTCTCCAATGCCTTTGTGATTGCCACAGTGTACCGGACCCGGAAACTGCACAC
T T L S N A F V I A T V Y R T R K L H T
250 270 290
CCCGGCTAACTACCTGATCGCCTCTCTGGACGTCACCGACCTGCTTGTGTCCATCCTGGT
P A N Y L I A S L D V T D L L V S I L V
310 330 350
GATACCCATCAGCACCATGTACACTGTACCGACCGCTGGACACTAAGCCAGGTGGTCTG
I P I S T M Y T V T D R W T L S Q V V C
370 390 410
TGACTTCTGGCTGTTCATCGGACATCACTTGTGCACTGCCTCCATCCTGCACCTCTGTGT
D F W L S S D I T C C T A S I L H L C V
430 450 470
CATCGCCCTGGACCGCTACTGGGCCATCACGGACGCCGTGGAGTACTCAGCTAAAAGGAC
I A L D R Y W A I T D A V E Y S A K R T
490 510 530
TCCCAAGAGGGCGGGCGGTTCATGATCGCGCTGGTGTGGGTCTTCTCCATCTCTATCTCGCT
P K R A A V M I A L V W V F S I S I S L
550 570 590
GCCGCCCTTCTTCTGGCGTCAGGCTAAGGCCGAAGAGGAGGTGTGCGGAATGCGTGGTGAA
P P F F W R Q A K A E E E V S E C V V N
610 630 650
CACCGACCACATCCTCTACACTGTCTACTCCACGGTGGGTGCTTTCTACTTCCCCACCCT
T D H I L Y T V Y S T V G A F Y F P T L
670 690 710
GCTCCTCATCGCCCTCTATGGCCGCATCTACGTAGAAGCCCGCTCCCGGATTTTGAAACA
L L I A L Y G R I Y V E A R S R I L K Q

Figure 4 (cont'd)

730 750 770
GACGCCCAACAGGACCGGCAAGCGCTTGACCCGAGCCCAGCTGATAACCGACTCCCCCGG
T P N R T G K R L T R A Q L I T D S P G

790 810 830
GTCCACGTCCTCGGTACCTCTATTAAGTTCGCGGGTTCCCGACGTGCCAGCGAATCCGG
S T S S V T S I N S R V P D V P S E S G

850 870 890
ATCTCCTGTGTATGTGAACCAAGTCAAAGTGCGAGTCTCCGACGCCCTGCTGGAAAAGAA
S P V Y V N Q V K V R V S D A L L E K K

910 930 950
GAAACTCATGGCCGCTAGGGAGCGCAAAGCCACCAAGACCCTAGGGATCATTGTTGGGAGC
K L M A A R E R K A T K T L G I I L G A

970 990 1010
CTTTATTGTGTGTTGGCTACCCTTCTTCATCATCTCCCTAGTGATGCCTATCTGCAAAGA
F I V C W L P F F I I S L V M P I C K D

1030 1050 1070
TGCCTGCTGGTTCCACCTAGCCATCTTTGACTTCTTCACATGGCTGGGCTATCTCAACTC
A C W F H L A I F D F F T W L G Y L N S

1090 1110 1130
CCTCATCAACCCCATATCTATACCATGTCCAATGAGGACTTTAAACAAGCATTCCATAA
L I N P I I Y T M S N E D F K Q A F H K

1150 1170 1190
ACTGATACGTTTAAAGTGCAAGTTGACTTGCCGTTTGCAGTGGGTCCCTAAGCGACCTT
L I R L S A Q V D L P F A V G P

1210 1230 1250
TGGGGACCAAGTTGTGTCTGCTTCCACAGGTAGGTCTGAATCTTCTTTCGCGGTTTCTGGG

1270 1290 1310
TCCCAGCGAGGCTCTCTCTCTGCGGCAAGGGCAATGGATCCTGAGAAGCCAGAATAGTCC

1330 1350 1370
TGAGAGAGAGCTCTGAAGGAGAAGTGTTGAAACCAAATGTAGAGCTTCCCTGCCCAGGAG

1390 1410 1430
GAGGCTCACTTCCCTCCCCTCAAGCCCCGGGCTCAGCACTGACCTGCGGCAGCCAATCCAA


1450 1470 1490
GGGGGTTGCAACTTTTAAAAATTGATATGGAAGGGAATCCCTGCCCTGCTTTGGTATCGT

1510 1530 1550
GATAATGCCCCACTAGAAGCAGTGTAATTGTTGTCTGAAGCCTGTCTGAGACAGAT

7/10

FIGURE 5

Peptide 8-30-84



○ RDC4 Differences

SUBSTITUTE SHEET

8/10

FIGURE 6

Consensus	M.....	..sp.nqs.e	gl...easn.s	lnatet.eaw	.p.tlqalk.	50
Pep 8-30-84	M.....	..SPLNQSAE	GLPQEASNRS	LNATETSEAW	DPRTLQALKI	
Pep 11	HEEPGAQCAP	PAPAGSETWV	PQANLSSAPS	QNCsAKDYIY	QDSISLPWKV	
RDC4 dog	M.....	..SPPNQSLZ	GLLQEASNRS	LNATETPEAW	GPETLQALKI	
5HT _{1A} hum	M.....DVLSPG	QGNTTSPPA	PFETGGNTTG	ISDVTVSQV	
TM Region*	
Consensus	sl.lLLs.lt	latvLnAFV	1.ti.ltrkL	htpANYLIGs	LavTDLLVSI	100
Pep 8-30-84	SLPVLLSVIT	LATVLSNAFV	LTTILLTRKL	HTPANYLIGS	LATTDLLVSI	
Pep 11	LLVHLLALIT	LATTLSNAFV	IATVYRTRKL	HTPANYLIAS	LDVTDLLVSI	
RDC4 dog	SLALLLSIIT	MATALSNAFV	LTTIFLTRKL	HTPANYLIGS	LAMTDLLVSI	
5HT _{1A} hum	ITSLLLGTLI	FCAVLGNACV	VAAIALERSL	QNVANYLIGS	LAVTDLMVSV	
TM Region	*****..I..	*****	*****	*****	*****..II	
Consensus	LvMpiS.ayt	vt.twt.gq.	lCDiwlssDI	tcCTaSiLHL	CvIALDRYWA	150
Pep 8-30-84	LVMPISMAyT	lTHWNFGQI	LCDIWLSSDI	TCCTASiLHL	CVIALDRYWA	
Pep 11	LVIPISThyT	VTDRWTLsQV	VCDFWLSSDI	TCCTASiLHL	CVIALDRYWA	
RDC4 dog	LVMPISIAyT	TTRTWsFGQI	LCDIWLSSDI	TCCTASiLHL	CVIALDRYWA	
5HT _{1A} hum	LVLPMaALyQ	VLNkWTlGQV	TCDLFIAlDV	LCCTSSiLHL	CAIALDRYWA	
TM Region	*****	*****	***.III.	*****	

9/10

FIGURE 6

151
ITDaleYsk. RT.graAVMI alvWvisICI SIPPf.WRQ akaqe.msdc 200
ITDALEYSKR RTAGHAATMI AIVWAISICI SIPPf.WRQ EKAQEEHSDC
ITDAVEYSK RPKRAAVMI ALWVFSISI SLPPF.WRQ AKAEVEVSEC
ITDALEYSKR RTAGRAAVMI ATWVISICI SIPPf.WRQ AKAQEDMSDC
ITDPIDYVNK RTPRAL.I SLTWLIGFLI SIPPMLGWT PEDRSDPDAC
..... **** IV... ****
.....

Consensus
Pep 8-30-84
Pep 11
RDC4 dog
5HT_{1A} hum
TM Region

201
.vnt...isYT IYSTCGAFYI Ps.LlilLYG RIY.aARNRI l.ppsl..gk 250
LVNTSQISYT IYSTCGAFYI PSVLLIILYG RIYRAARNRI LNPPSL.GK
VVNTDHILYT VYSTVGAFYI PTLILLIILYG RIYVEARSRI LKQTPNRTGK
QVNTSQISYT IYSTCGAFYI PSVLLIILYG RIYVAARNRI LNPPSLY.GK
TISKDH.GYT IYSTFGAFYI PLLLMLVLYG RIFRAARFRI RKTvKKVE.K
..... **** V. ****
.....

Consensus
Pep 8-30-84
Pep 11
RDC4 dog
5HT_{1A} hum
TM Region

251
rfttaqlitg S.gs...s...sg.r..a ..p..f..... .k....s...l. 300
RFTTAHLITG SAGSVCSLNS SLHEGHS HA GSPLFFNHVK IKLADSALER
RLTRAQLITD SPGSTSSVTS INSRVPDVPs ESGSPVYVQ VKVRVSDALL
RFTTAQLITG SAGSSLCSLS PSLQEERSHA AGPPLFFNHV QVKLAEGVLE
TGADTRH.GA SPAQPKKSV NGESGSRNWR LGVESKAGGA LCANGAVRQ
.....

Consensus
Pep 8-30-84
Pep 11
RDC4 dog
5HT_{1A} hum
TM Region

SUBSTITUTE SHEET

FIGURE 6

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application

No. PCT/US91/03200

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶
According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): C07H 21/00; C12Q 1/68

US Cl: 536/27; 435/6

II. FIELDS SEARCHED

Minimum Documentation Searched ⁷	
Classification System	Classification Symbols
US	435/6, 29, 34, 69.1, 70.3, 71.2, 91, 172.3, 240.2, 320.1, 820; 436/501, 503, 86, 811; 536/27; 935/4, 9, 19, 29, 34, 36, 41, 70, 72, 77, 78, 88
Documentation Searched other than Minimum Documentation to the extent that such documents are included in the fields searched ⁸	

III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹

Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	Proceedings of the National Academy of Sciences (USA), Volume 80 issued November 1983, Zabel et al., "High Resolution chromosomal localization of human genes for amylase, proopiomelanocortin, somatostatin, and a DNA fragment (D3S1) by <u>in situ</u> hybridization", pages 6932-6936, see especially Figure 1 on page 6933.	1, 2, 28-30, 32-34, 77
X	SCIENCE, Volume 244, issued 05 May 1989, Libert et al., "Selective Amplification and Cloning of Four New Members of the G Protein-Coupled Receptor Family", pages 569-572, see especially the legend of Figure 1 on page 570.	1-4, 8, 9, 28-30, 32-34, 77
Y	SCIENCE, Volume 241, issued 29 July 1988, Julius et al., "Molecular Characterization of a Functional cDNA Encoding the Serotonin 1c Receptor", pages 558-564, see especially the abstract.	1-4, 8-22, 24-26, 28-30, 32-35, 77, 83

* Special categories of cited documents: ¹⁰

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

26 August 1991

Date of Mailing of this International Search Report

09 SEP 1991

International Searching Authority

ISA/US

Signature of Authorized Officer

ARDIN MARSCHER

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	WO, A, 89/08149 (AXEL ET AL.) 08 September 1989, see especially the abstract.	1-4, 8-22, 24-26, 28-30, 32-35, 77, 83
Y	Proceedings of the National Academy of Sciences (USA), Volume 84, issued June 1987, Lubbert et al., "cDNA cloning of a serotonin 5-HT _{1c} receptor by electrophysiological assays of mRNA-injected Xenopus oocytes", pages 4332-4336, see especially the abstract.	1-4, 8-22, 24-26, 28-30, 32-35, 77, 83
Y	NATURE, volume 335, issued 22 September 1988, Fargin et al., "The Genomic clone G-21 which resemble a α -adrenergic receptor sequence encodes the 5-HT _{1A} receptor", pages 358-360, see especially the abstract.	1-4, 8-22, 24-26, 28-30, 32-35, 77, 83
Y	The Journal of Biological Chemistry, Volume 265, No. 10, issued 05 April 1990, Albert et al., "Cloning, Functional Expression, and mRNA Tissue Distribution of the Rat 5-Hydroxytryptamine _{1A} Receptor Gene", pages 5825-5832, see especially the abstract.	1-4, 8-22, 24-26, 28-30, 32-35, 77, 83

Form PCT/ISA/210 (extra sheet) (Rev.11-87)

FURTHER INFORMATION

CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter ¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out ¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING¹

This International Searching Authority found multiple inventions in this international application as follows:

See attached sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
1-4, 8-22, 24-26, 28-30, 32-35, 77, 83 telephone practice
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
☐ No protest accompanied the payment of additional search fees.

ATTACHMENT A (CONTINUATION OF PART VI)

OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

I. Claims 1-4, 8-22, 24-26, 28-30, 32-35, 77, and 83, drawn to human 5-HT_{1A} receptor nucleic acid and a binding assay using cells containing this nucleic acid, classified in Class 435, subclass 6.

II. Claims 5-7, drawn to human 5-HT_{1A} receptor protein, classified in Class 530, subclass 399.

III. Claim 23, drawn to ligands that bind to said receptor protein, classified in Class 530, subclass 23.

IV. Claim 27, drawn to pharmaceutical compositions that bind to said receptor protein, classified in Class 514, subclass 43.

V. Claims 31, 73, and 74, drawn to a hybridization detection method, classified in Class 435, subclass 6.

VI. Claims 36-39, drawn to a pharmaceutical composition that binds to mRNA, classified in Class 514, subclass 44.

VII. Claims 40-46, drawn to a mRNA treatment method, classified in Class 435, subclass 91.

VIII. Claims 47-49, drawn to an antibody, classified in Class 530, subclass 387.

IX. Claims 50, 68, and 71, drawn to an antibody pharmaceutical composition, classified in Class 514, subclass 21.

X. Claims 51-57, 69, and 72, drawn to an antibody treatment method, classified in Class 435, subclass 70.1.

XI. Claim 58, drawn to receptor detection via antibody, classified in Class 435, subclass 7.21.

XII. Claims 59-64, 78, 79, and 82, drawn to transgenic animals, classified in Class 435, subclass 240.1.

XIII. Claims 65-67, 70, 80, and 81, drawn to assays using transgenic animals, classified in Class 435, subclass 6.

XIV. Claims 75 and 76, drawn to a method of preparation of receptor protein, classified in Class 435, subclass 69.1.

Group I is directed to a nucleic acid and its first method of use claimed which is a binding assay using cells containing said nucleic acid. The remainder of the Groups are either further methods of use or clearly distinct compositions such as proteins, antibodies, etc., and their respective methods of use all requiring separate search due to either their different classification or clearly distinct subject matter.

This Page Blank (uspto)